



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/56		A1	(11) International Publication Number: WO 94/28904
			(43) International Publication Date: 22 December 1994 (22.12.94)
(21) International Application Number: PCT/US93/09349		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 September 1993 (28.09.93)			
(30) Priority Data: 08/077,359 15 June 1993 (15.06.93) US			
(71) Applicant: PHERIN CORPORATION [US/US]; Suite 240, 535 Middlefield Road, Menlo Park, CA 94025 (US).			
(72) Inventors: BERLINER, David, L.; 380 Selby Lane, Atherton, CA 94027 (US). ADAMS, Nathan, William; 4858 South Rockhill Lane, Salt Lake City, UT 84123 (US). JENNINGS-WHITE, Clive, L.; 3096 South 2300 East, Salt Lake City, UT 84109 (US).			
(74) Agents: SUYAT, Reginald, J. et al.; Heller, Ehrman, White & McAuliffe, 333 Bush Street, San Francisco, CA 94104 (US).		Published With international search report. With a request for rectification under Rule 91.1(f).	

(54) Title: ANDROSTANE STEROIDS AS NEUROCHEMICAL INITIATORS OF CHANGE IN HUMAN HYPOTHALAMIC FUNCTION AND RELATED PHARMACEUTICAL COMPOSITIONS AND METHODS

(57) Abstract

The invention relates to a method of altering hypothalamic function in an individual. The method comprises nasally administering a human semiochemical, e.g. an Androstane steroid, or a pharmaceutical composition containing a semiochemical, such that the ligand semiochemical binds to a specific neuroepithelial receptor. The steroid or steroids is/are preferably administered in the form of a pharmaceutical composition containing one or more pharmaceutically acceptable carriers. Other embodiments of the invention include pharmaceutical compositions containing the steroids.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ANDROSTANE STEROIDS AS NEUROCHEMICAL INITIATORS OF
CHANGE IN HUMAN HYPOTHALAMIC FUNCTION AND
RELATED PHARMACEUTICAL COMPOSITIONS AND METHODS

BACKGROUND

5 Technical Field

This invention relates generally to pharmaceutical compositions and methods for effectuating change in human hypothalamic function, thereby altering certain behavior and physiology
10 mediated by the hypothalamus of individuals. More particularly, the invention relates to the use of certain Androstene steroids as neurochemical effectuators of physiology and behavior.

Description of the Related Art

15 The present invention relates to certain compounds, namely Androstane steroids, particularly Androstene steroids and related compounds as will be described herein, and methods of using these
20 compounds as human semiochemicals in order to alter hypothalamic function, thereby affecting certain consequent behavior and physiology, e.g., the reduction of anxiety. Androstane steroids are typified by testosterone and they are characterized by a four ring steroidal structure, a methylation at

-2-

the 13-position and at the 10-position. Androstenes are a subset of Androstanes and have at least one double bond. Ohloff, G. et al. (Helv. Chim. Acta (1983) 66:192-217), which is incorporated herein by reference, have shown that several members of this group of steroids have an odor which varies with different isomeric, diastereomeric, and enantiomeric forms. Some members of this group have been reported to act as a pheromone in some mammalian species - for instance, 5 α -androst-16-en-3-one and 5 α -androst-16-en-3 α -ol in pigs (Melrose, D.R., et al., Br. vet. J. (1971) 127:497-502). These 16-Androstenes produced by the boar induce mating behavior in estrus sows (Claus, et al., Experimentia (1979) 35:1674-1675).

Some studies have noted that, in some species, various characteristics of certain 16-Androstenes (including 5 α -Androst-16-en-3 α 2-ol and 5 α -Androst-16-en-3-one), such as concentration, metabolism, and localization, are sexually dimorphic (Brooksbank et al., J. Endocr. (1972) 52: 239-251; Claus, et al., J. Endocr. (1976) 68:483-484; Kwan, et al., Med. Sci. Res. (1987) 15:1443-1444). For instance, 5 α -Androst-16-en-3 α -ol and 5 α -Androst-16-en-3-one, as well as Androsta-4,16-dien-3-one, have been found at different concentrations in the peripheral blood, saliva and axillary secretions of men and of women (Kwan, T.K., et al., Med. Sci. Res. (1987) 15:1443-1444), and their function as a human pheromone, to the extent of affecting choice and judgement, has been suggested (Id.; see also Gower, et al., "The Significance of Odorous Steroids in Axillary Odour", In, Perfumery, pp. 68-72, Van Toller and Dodd, Eds., Chapman and Hall, 1988); Kirk-Smith, D.A., et al., Res. Comm. Psychol. Psychiat. Behav. (1978) 3:379). Androstenol (5 α -androst-16-en-3 α -ol)

-3-

has been claimed to exhibit a pheromone-like activity in a commercial men's cologne and women's perfume (Andron™ for men and Andron™ for women by Jövan). Japanese Kokai No. 2295916, refers to perfume compositions containing androstenol and/or its analogues. 5 α -Androstadien-3 β -ol (and perhaps the 3 α -ol) has also been identified in human axillary secretion (Gower, et al., Supra at 57-60. On the other hand, there is little agreement in the literature as to whether or not any putative pheromone actually plays any role in the sexual or reproductive behavior of mammals, particularly of humans. See: Beauchamp, G.K., et al., "The Pheromone Concept in Mammalian Chemical Communication: A Critique", In: Mammalian Olfaction, Reproductive Processes and Behavior, Doty, R.L., Ed., Academic Press, 1976). See also: Gower, et al., supra at 68-73.

An embodiment of the subject invention concerns the non-systemic, nasal administration of certain Androstane and Androstene steroids to affect a specific behavioral or physiological response in human subjects, e.g., a reduction of negative affect, mood, and character traits. In particular, nasal administration provides for contacting neurochemical receptors of a heretofore poorly understood neuroendocrine structure, commonly known as the vomeronasal organ ("VNO"; also known as "Jacobson's organ"), with one or more steroid(s) or with compositions containing the steroid(s). This organ is accessed through the nostrils of most higher animals - from snakes to humans, and has been associated, inter alia, with pheromone reception in certain species (see generally Muller-Schwarze & Silverstein, Chemical Signals, Plenum Press, New York

-4-

(1980)). The axons of the neuroepithelia of the vomeronasal organ, located supra palatinal, form the vomeronasal nerve and have direct synaptic connection to the accessory olfactory bulb and indirect input
5 from there to the cortico-medial amygdaloid basal forebrain and hypothalamic nuclei of the brain. The distal axons of terminalis nerve neurons may also serve as neurochemical receptors in the VNO. Stensaas, L.J., et al., J. Steroid Biochem. and
10 Molec. Biol. (1991) 39:553. This nerve has direct synaptic connection with the hypothalamus.

Johnson, A. et al. (J. Otolaryngology (1985) 14:71-79) report evidence for the presence of the vomeronasal organ in most adult humans, but
15 conclude that the organ is probably non-functional. Contravening results which suggest that the VNO is a functional chemosensory receptor are reported by Stensaas, L., et al., supra; and by Moran, D.T., et al., Garcia-Velasco, J. and M. Mondragon; Monti-
20 Bloch, L. and B. Grosser all in J. Steroid Biochem. and Molec. Biol. (1991) 39.

It is apparent that it would be desirable to identify and synthesize human semiochemicals and pheromones and to develop pharmaceutical compositions
25 and methods of use to influence hypothalamic function. This invention relates to the unexpected discovery that, when nasally administered to human subjects, certain neurochemical ligands, particularly Androstane steroids, more particularly Androstene
30 steroids and related compounds, or pharmaceutical compositions containing Androstanes, Androstenes or related compounds, specifically bind to chemoreceptors of certain nasal neuroepithelial cells and this binding generates a series of
35 neurophysiological responses resulting in an

-5-

alteration of hypothalamic function of an individual. When properly administered, the effect of certain of these compounds on the hypothalamus affects the function of the autonomic nervous system and a variety of behavioral-or physiological phenomena which include, but are not limited to the following: anxiety, premenstrual stress, fear, aggression, hunger, blood pressure, and other behavioral and physiological functions normally regulated by the hypothalamus. Otto Appenzeller. The Autonomic Nervous System. An introduction of basic and clinical concepts (1990); Korner, P.I. Central nervous control of autonomic cardiovascular function, and Levy, N.M. and Martin, P.J. Neural control of the heart, both in Handbook of Physiology; Section 2: Cardiovascular System - the heart, Vol I, Washington DC, 1979, American Physiological Society; Fishman, A.P., et al. editors, Handbook of Physiology. Section 3: Respiratory System. Vol. II. Control of breathing. Bethesda MD. 1986. American Physiological Society.

In some instances a single Androstane steroid, or related compound, is administered, in some instances combinations of Androstane steroids and/or related compounds are administered and in some instances one or more Androstane steroids are co-administered along with one or more Estrane or Estrene steroids or a related compound.

SUMMARY OF THE INVENTION

Accordingly, it is an object of this invention to provide pharmaceutical compositions which contain human semiochemicals or pheromones and are suitable for nasal administration in an individual.

-6-

It is also an object of this invention to provide methods of using these compositions to alter hypothalamic function of an individual.


It is a further object of this invention to
5 provide methods of using these compositions to affect physiological and behavioral functions of individuals which are normally regulated by the hypothalamus.

Finally, it is an object of this invention to provide methods of altering hypothalamic function
10 which have the following advantages: 1) administration directly to the chemoreceptors in the nasal passage and the vomeronasal organ, without pills or needles - i.e., non-invasively; 2) a mode of drug action through the nervous system and not
15 through the circulatory system - thus brain function can be affected without consideration of the blood-brain barrier; 3) a direct means of affecting the hypothalamus - there is only one synaptic junction between pheromone receptors and the hypothalamus;
20 and, 4) providing a highly specific drug effect, thereby greatly reducing the potential for undesirable side-effects - this because sensory nerves are addressed to a specific location in the brain.

25 Additional objects, advantages and novel features of the invention will be set forth in part in the description which follows, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by
30 practice of the invention.

Objects of this invention are achieved by providing a pharmaceutical composition suitable for nasal administration in an individual. The composition contains a pharmaceutically acceptable
35 carrier and an Androstane steroid with the formula:

-7-



wherein P_1 is selected from the group consisting of
 oxo, α -(β -) hydroxy, α -(β -) acetoxy, α -(β -)
 propionoxy, α -(β -) methoxy, α -(β -) lower acyloxy, α -
 (β -) lower alkyloxy, and α -(β -) benzoyloxy; P_2 is
 5 selected from the group consisting of methyl,
 hydroxymethyl, acyloxymethyl, alkoxymethyl, lower
 alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl;
 P_3 is absent or is selected from the group consisting
 of methyl, hydroxymethyl, acyloxymethyl,
 10 alkoxymethyl, lower alkyl, hydroxyalkyl,
 acyloxyalkyl, and alkoxylalkyl; P_4 is selected from
 the group consisting of hydrogen, oxo, halo, hydroxy,
 alkoxy, and acyloxy; P_5 represents one or 2
 substituents, wherein P_5 comprises one or two
 15 hydrogen atoms, methyl, methylene, or one or two halo
 atoms; P_6 is hydrogen or halo; and "a", "b", "c",
 "d", "e", "f", and "h" are alternative sites for
 optional double bonds.

One class of preferred steroids has "b" as
 20 a double bond, particularly wherein "c" or "d" is
 also a double bond. Another preferred class has "a"
 and "c" as double bonds. Yet another preferred class
 contains P_3 as a methyl group, "h" as an optional
 double bond, and P_5 as methylene or one or two

-8-

hydrogen atoms. A class of steroids wherein "a" or "b" is a double bond is also preferred.

By halo, it is meant, F, Cl, Br, or I. The term lower alkyl, lower alkoxy, etc., is meant to encompass carbon chains of 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms.

Other objects of this invention are achieved by providing a method of altering hypothalamic function and/or autonomic function in an individual. A ligand for a chemoreceptor displayed on the surface of a nasal neuroepithelial cell is provided wherein the cell is a part of tissue other than olfactory epithelia; and, the ligand is administered within a nasal passage of the individual such that the ligand binds specifically to the chemoreceptor, resulting in an alteration of hypothalamic function of the individual.

All embodiments of this application relate to and include the functional equivalents of the steroid structures disclosed in these embodiments and to those modified steroids which demonstrate said functional equivalence, whether or not the modified steroids are explicitly disclosed.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the synthesis of Androsta-4,16-dien-3-one, Androsta-4,16-dien-3 α -ol, and Androsta-4,16-dien-3 β -ol.

Figure 2 illustrates the synthesis of Androsta-5,16-dien-3 α -ol and Androsta-5,16-dien-3 β -ol.

Figure 3 illustrates an alternate synthesis of Androsta-4,16-dien-3-one.

Figure 4 is a graphic representation of the electrophysiological effect on receptor potential of

-9-

the localized administration of particular steroids to the vomeronasal organ of female subjects (4A) and to the olfactory epithelium (4C). Figure 4B is a graphic comparison of the effect of an Androstane on the VNO receptor potential of male and female subjects.

Figure 5 is a graphic representation of the electrophysiological effect of the localized administration of particular steroids to the vomeronasal organ of male (5A) and female (5B) subjects.

Figure 6 depicts various autonomic responses of female subjects to an Androstane. A = receptor potential of the vomeronasal neuroepithelium; B = change in cortical alpha activity of an electroencephalogram (%); C = change in galvanic skin response (K-ohms); D = change in peripheral arterial pulse (counts/min.); E = change in skin temperature (degrees C.); and, F = change in respiratory frequency (counts/min.).

Figure 7 depicts changes in receptor potential of the VNO after exposure of 5 females to two different Androstanes.

Figure 8 depicts sexual dimorphism in local and autonomic responses to the stimulation of the VNO with vomeropherins. Various vomeropherins (200 fmoles) and the diluent control were administered to 30 male and 30 female subjects (ages 20 to 45) as described. Bars indicate the mean response of the population.

Figs. 8A & B: EVG responses were measured as described in male (A) and female (B) subjects.

-10-

Figs. 8C & D: Electrodermal activity was measured as described. Changes (measured in $\times\Omega$) in response due to delivery of vomeropherins to the VNO of each subject are shown in male (C) and female (D) subjects.

Figs. 8E & F: Alpha-cortical activity was measured as described. Changes in response due to delivery of vomeropherins to the VNO of male (E) and female (F) subjects.

10 Figs. 8G & H: Skin temperature (ST) was measured as described. Changes in response due to delivery of vomeropherins to the VNO of each subject are shown in male (G) and female (H) subjects.

The Compounds in the graphs are:

- 15 A = 1, 3, 5(10),16-Estratetraen-3-yl acetate
B = Androsta-4,16-dien-3-one
C = 1,3,5(10),16-Estratetraen-3-ol
D = 3-Methoxy-Estra-1,3,5(10),16-tetraene
E = Androsta-4,16-dien-3 α -ol
20 F = Androsta- 4,16-dien-3 β -ol

Figure 9 depicts electro-olfactograms of male and female subjects induced by stimulation of the OE with olfactants and vomeropherins A: 400 fmoles of the olfactants 1-carvone and cineole as well as 200 fmoles of the vomeropherins A, B, C, D and F; and the stereoisomer E were applied separately as one second pulses to the OE of 20 subjects (both male and female) and each EOG response was recorded as described. The olfactants as well as E and B produced significant ($p < 0.01$) local response. B: 400 fmoles of the olfactants 1-carvone and cineole do not

-11-

induce a significant EVG response when delivered to the VNO of male and female subjects.

Figure 10 depicts the electrophysiological effect of the following vomeropherins on the
5 vomeronasal organ of 20 female subjects:

G = Androst-4-en-3-one

H = Androsta-4,16-diene-3,6-dione

J = 10,17-Dimethylgona-4,13(17)-dien-3-one

K = 1,3,5(10),16-Estratetraen-3-ol-methyl ether

10 L = 1,3,5(10),16-Estratetraen-3-yl-propionate

EVG = Electro-vomeronasogram

GSR = Galvanic Skin Response

= Electrodermal Activity, EDA

ST = Skin Temperature

15 Figure 11 depicts the electrophysiological effect of vomeropherins on the vomeronasal organ of 20 male subjects.

M = 1,3,5(10)-Estratrien-3-ol

DETAILED DESCRIPTION OF THE INVENTION

20 I. Definitions

An "affect" is a transient feeling state. Typical negative affects are feelings of nervousness, tenseness, shame, anxiousness, irritability, anger, rage, and the like. "Moods" are longer lasting
25 feeling states such as guilt, sadness, hopelessness, worthlessness, remorsefulness, misery, unhappiness and the like. "Character traits" are more permanent aspects of an individual's personality. Typical negative character traits are sensitivity,
30 regretfulness, blameworthiness, stubbornness,

-12-

resentfulness, bitterness, timidity, laziness and the like.

"Androstane steroids" are aliphatic polycyclic hydrocarbons characterized by a four-ring steroidal structure with a methylation at the 10- and 13-positions. An Androstene is a subset of Androstanes commonly understood to mean that the compound has at least one double bond. Commonly, unless a compound is described as a gonane, it is understood that the compound has an 18- carbon group. However, it is intended herein that 18-Nor-Androstanes are herein regarded as 16-Androstane steroids. Furthermore, all derivatives which have the structural characteristics described above are also referred to generically as Androstane steroids.


A "chemoreceptor" is a receptor molecule displayed on the surface of a "chemosensory" neuroepithelial cell which binds in a stereospecific fashion to a particular ligand or ligands. This specific binding initiates a signal transduction which initiates an afferent nerve impulse. Chemoreceptors are found, inter alia, in taste buds, olfactory epithelium and vomeronasal tissue.

"Estrene steroids", as the term is used herein, are aliphatic polycyclic hydrocarbons with a four-ring steroidal structure, at least one double bond in the A-ring, no methylation at the 10-position and an oxo, hydroxyl or hydroxyl derivative such as an alkoxy, ester, benzoate, cypionate, sulfate or glucuronide, at the 3-position. Derivatives which contain these structural characteristics are also referred to generically as Estrene steroids.

The following structure shows the four-ring steroidal structure common to Androstane and Estrene steroids. In describing the location of groups and

-13-

substituents, the following numbering system will be employed:



"Sexually dimorphic" refers to a difference in the effect of, or response to, a pharmaceutical agent between males and females of the same species.

An "effective amount" of a drug is a range of quantity and/or concentration which brings about a desired physiological and/or psychological effect when administered to an individual in need of the drug. In the present case, a needy individual is one with a physiological or behavioral trait which is normally regulated by the hypothalamus and wherein it is desirable to affect the function of the hypothalamus or the trait. The effective amount of a given drug may vary depending upon the function to be affected, the desired effect, route of administration, and the like. For example, when the steroid is administered as a solution applied to the facial skin of a subject an effective concentration is from 1 microgram/ml to 100 $\mu\text{g/ml}$, preferably 10 to 50 $\mu\text{g/ml}$ and most preferably 20 to 30 $\mu\text{g/ml}$. When the steroid is introduced directly into the VNO an effective amount is about 1 picogram to about 1 nanogram, more preferably about 10 picograms to about 50 picograms. When the steroid is administered to the nasal passage, by ointment, cream or aerosol, or the like, an effective amount is about 100 pg to about 100 micrograms, preferably about 1 ng to about

-14-

10 micrograms. It follows that some drugs may be effective when administered by some routes, but not effective when administered by other routes.

The "hypothalamus" is the portion of the diencephalon comprising the ventral wall of the third ventricle below the hypothalamic sulcus and including structures forming the ventricle floor, including the optic chiasma, tuber cinereum, infundibulum, and mammillary bodies. The hypothalamus regulates the autonomic nervous system and controls several physiological and behavioral functions such as the so-called fight and flight responses, sexual motivation, water balance, sugar and fat metabolism, hunger, regulation of body temperature, endocrine secretions, and others. The hypothalamus is also the source of vasopressin which regulates blood pressure, and oxytocin which induces parturition and milk release. All hypothalamic functions are potentially modulatable by the semiochemical therapy described herein.

A "ligand", as used herein, is a molecule which acts as a chemical signal by specifically binding to a receptor molecule displayed on the surface of a receptor cell, thereby initiating a signal transduction across the cell surface. Binding of ligands to chemosensory receptors can be measured. Chemosensory tissue, such as vomeronasal neuroepithelium or olfactory neuroepithelium, contains a multiplicity of neuroreceptors cells, each displaying at least one cell surface receptor. Many of the receptor molecules have identical ligand specificity. Therefore, when the tissue is exposed to a ligand for which it has specificity (for example a exposure of the VNO to a semiochemical) a summated

-15-

change in cell surface receptor potential can be measured.

As used herein, "lower alkyl" means a branched or unbranched saturated hydrocarbon chain of 1 to 4 carbons, such as, for example, methyl, ethyl, n-propyl, i-butyl and the like. "Alkoxy" as used herein is used in its conventional sense to mean the group -OR wherein R is alkyl as herein defined.

A "pheromone" is a substance that provides chemical means of communication between members of the same species through secretion and nasus reception. In mammals pheromones are usually detected by receptors in the vomeronasal organ of the nose. Commonly, pheromones effect development, reproduction and related behaviors. A "semiochemical" is a more general term which includes pheromones and describes a substance from any source which functions as a chemosensory messenger, binds to a specific neuroepithelial receptor, and induces a physiological or behavioral effect. A "vomeropherin" is a semiochemical whose physiologic effect is mediated through the vomeronasal organ.

A picogram (pg) is equal to .001 nanograms (ng). A ng is equal to .001 micrograms (μ g). A μ g is equal to .001 mg.

II. Modes for Carrying Out the Invention

A. Androstanes useful in the Invention

The invention is directed to a group of certain Androstane steroids. Testosterone (17-hydroxy-Androsta-4-en-3-one) is a typical Androstane.

Androstanes especially suitable for use in the present invention include those where, independently, P_1 = oxo, α -hydroxy, β -hydroxy; P_2 =

-16-

methyl, lower alkyl, hydroxymethyl, hydroxyalkyl; P_3 = hydrogen or methyl; P_4 = hydrogen, hydroxy, or oxo; P_5 = hydrogen or methyl; and there is at least one double bond, usually at the 4- or 16- position.













































- 5 Preferred Androstanes include Androsta-4,16-dien-3-one (P_1 = oxo, a = double bond, P_2 = methyl, P_3 , P_4 , P_5 = hydrogen, commercially available from Steraloids, Inc.), Androsta-4,16-dien-3 β -ol (P_1 = β -OH, a = double bond, P_2 = methyl, P_3 , P_4 , P_5 = hydrogen), and 6-keto-Androsta-4,16-diene-3-one (P_1 = oxo, a = double bond, P_2 = methyl, P_3 , P_5 = hydrogen, P_4 = oxo), synthesis of which are described herein.
- 10

- A subset of Androstanes within the group are believed to be novel. Syntheses are described herein for the following compounds as designated on the chart: 17-methylene-Androst-4-en-3 β -ol (A3/N3), 17-methylene-Androst-4-en-3 α -ol (A4/N3), 17-methylene-6-oxo-Androst-4-en-3-one (A6/N3), and 6 β -OH-Androsta-4,16-dien-3-one (A11/N1).
- 15

- 20 Chart 1 includes androstanes to which the invention is directed, but do not limit its scope. The synthesis diagrams that follow depict intermediate and substructure syntheses for the preparation of these androstanes:

-17-

CHART I: ANDROSTRANES


A \ N	1	2	3	4
1	 KNOWN	 KNOWN	 KNOWN	 KNOWN
2	 KNOWN	 KNOWN	 KNOWN	
3	 KNOWN	 KNOWN	 KNOWN	
4	 KNOWN	 KNOWN	 KNOWN	
5	 KNOWN	 KNOWN		
6	 KNOWN	 KNOWN		
7	 KNOWN			
8	 KNOWN	 KNOWN	 KNOWN	
9	 KNOWN	 KNOWN		
10	 KNOWN	 KNOWN		
11	 KNOWN	 KNOWN		

-18-


SUBSTRUCTURE SYNTHESSES

Referring to the preceding table, the following are exemplary syntheses for intermediates in a given row (A1 through A11) or column (N1 through N4).

5

A1: Type A



A2:





10 This is a commercially available substructure, for example, DEHYDRO EPI ANDROSTERONE.

-19-




A3:



(A1)





(A3)






(Michio Matsui and David K. Fukushima, J. Org. Chem., 1970, Vol. 35, No. 3, p. 561-564).


5 A4:



(A1)





48%



Ohloff, G. et al. (Helv. Chim. Acta (1983) 66: 192-217).


-20-

A5:





(A1)

CHLORANIL





(A5)


CHLORANIL
45%

German Off. 2,631,915.



A6:



(A2)

 $\text{CrO}_3/\text{H}_2\text{SO}_4$ 

(A6)



 $\text{CrO}_3/\text{H}_2\text{SO}_4$
36%

5



J. Römer, H. Wagner, and W. Sihade, Steroids, 1988, 51/5-6, p. 577-581).

-21-


A7:



Several steps





Several steps





(Habermehl, et al., Z. Naturforsch. (1980) 256: 191-195).


5 A8:



MeO \times OMe/H⁺





MeO \times OMe/H⁺
44%





-SEE Example 15-

-22-

A9:




DICHLORODICYANO BENZOQUINONE







DICHLORODICYANO BENZOQUINONE

43%




Ohloff, G. et al. (Helv. Chim. Acta (1983) 66: 192-217).

5 A10:

 LiAlH_4  LiAlH_4



30%




V. I. Mel'nikova and K. K. Pivnitskii, Zhurnal Organicheskoi Khimii, 1972, Vol. 8, No. 1, pp. 68-74).

-23-



A11:



(A8)



(A11)




-SEE EXAMPLE 19-


- 24 -

Type N

N1:



N2:



5


1. Robert H. Shapiro and Carl Djerassi, J. Am. Chem. Soc., 1964, 86, 2825.

2. Pilar Lupón, Frances C. Canals, Arsenio Iglesias, Joan C. Ferrer, Albert Palomar, and Juan-Julio Bonet, J. Org. Chem. 1988, 53, 2193-2198.


10

-25-


N3:



as in




and




1. Günther Drefahl, Kurt Ponold and Hans Schick, *Berichte*, 1965, 98, 604.

- 5 2. Richard H. Peters, David F. Crows, Mitchell A. Avery, Wesley K. M. Chong, and Masako Tanabe, *J. Med. Chem.*, 1989, 32, 1642.


N4:



as in



and




- 10 1. Franz Sondheimer, O. Mancera, M. Urquiza & G. Rosenkranz, *J. Am. Chem Soc.*, 1955, 77, 4145.

2. William F. Johns, *J. Org. Chem.*, 1961, 26, 4583.


-26-

Methylandrostenes

German Off. 2,631,915 teaches preparation of




with a methyl group at any one of the following
5 positions: 1 α , 2 α , 4, 6 α , 6 β , 7 α , and 16.




6-METHYLANDROSTA-4,6-DIEN-3-ONE

German Off. 2,428,679.

Synthesis of the 17-METHYLANDROSTENES:






and

















Daniel Bertin and Lucien Nedelac, Mémoires Présentées
10 a la Société Chimique, 1964, No. 345, p. 2140.

Synthesizable compounds therefore include these,
together with those derived from them; i.e., N1 with
methyl at 1 α , 2 α , 4, 6 α , 6 β , 7 α , 16 or 17 combined
with A1, A3, A4, A5, A8, A9, A10 or A11, as well as
15 A2 or A6 with a 17-methyl.

-27-


Haloandrostenes

U.S. Patent 3,413,321.



5 German Off. 2,631,915.

-28-



European Patent Application EP 208,497.

Synthesizable compounds therefore include these, together with those derived from them; i.e.,


- 5 (4-Chloro, 4-Bromo, 6 α -Chloro, 6 α -Bromo, 6 β -Chloro, 6 β -Bromo, or 6 β -Iodo)-A1 in combination with N1, N2, N3, or N4. In addition, (17-Fluoro, 17-Chloro, 17-Bromo, or 17-Iodo)-N1 in combination with A1, A2, A3, A4, A5, A6, A8, A9, A10 or A11.

10

B. Estrenes useful in the Invention

The invention is additionally directed to compositions and methods involving the combination of the aforementioned Androstane steroids with certain Estrene steroids which are structurally related to

15 Estradiol (also referred to as 1,3,5(10)-Estratriene-3,17 β -diol). Estrene steroids useful in this invention have the formula:



-29-

wherein R_1 is selected from the group consisting essentially of one or two hydrogen atoms, methyl, methylene, and one or two halo atoms; R_2 is absent or is selected from the group consisting essentially of hydrogen and methyl; R_3 is selected from the group consisting essentially of oxo, hydroxy, lower alkoxy, lower acyloxy, benzoyl, cypionyl, glucuronide and sulfonyl; R_4 is selected from the group consisting essentially of hydrogen, hydroxy, lower alkoxy, lower acyloxy and halo; R_5 is absent or is selected from the group consisting essentially of hydrogen, hydroxy, lower alkoxy and lower acyloxy; R_6 is a hydrogen or a halo; and "a" represents optional aromatic unsaturation of ring A of said steroid, or "b", "c", and "d" are each optional double bonds; and "e", "f", "g", "h", "i" and "j" are each optional double bonds. In this embodiment, the steroid is preferably administered in the form of a pharmaceutical composition containing one or more pharmaceutically acceptable carriers.

A preferred class of compounds are those in which "a" is present and "g", "h" or "i" are optional double bonds. The class wherein "h" and "i" are both double bonds is also preferred. Another preferred class contains "b", "c" or "j" as a double bond. Yet another class contains "c" and "d" as double bonds. Still another class contains R_2 as methyl and (1) "e" as a double bond, (2) R_1 is methylene or a single hydrogen, or (3) "f" is a double bond.

Preferred estrenes include 1,3,5(10)-Estratriene-3, 17 β -diol; 1,3,5(10)-Estratriene-3,16 α , 17 β -triol; 1,3,5(10)-Estratriene-3-ol-17-one; 1,3,5(10),16-Estratetraen-3-ol; 1,3,5 (10),16-Estratetraen-3-ol methyl ether; and 1,3,5(10),16-Estratetraen-3-yl acetate.

-30-

Most of these steroids and their glucuronide, sulfate, cypionate, and benzoate derivatives, are compounds known in the art and are commercially available, e.g., from Sigma Chemical Co., Aldrich Chemical Co., etc. Alkoxy derivatives and their syntheses are also known in the art and shown in U.S. Patent No. 2,984,677.

- 1,3,5(10),16-Estratetraen-3-ol is available from Research Plus, Inc. and from Steraloids, Inc.
- 10 Another synthesis of this compound, as well as syntheses of the acetate and propionate derivatives are described in the commonly assigned, co-pending continuation-in-part of U.S.S.N. 07/903,525, which is in turn a continuation-in-part of U.S.S.N.
- 15 07/707,862, incorporated herein by reference.

C. Synthetic Methods

1. Preparation of 3-, 5-, 6-, 18- and 19- position derivatives.

The compounds used in the methods of this invention are Androstane steroids substituted at the 3-, 5-, 6-, 18- and 19- positions. Many of the 3- and 5-substituted steroids are known compounds which may be derived from 17-hydroxy-and 17-oxo-steroids (commercially available e.g. from Aldrich Chemical Co) by elimination or reduction to the Δ^{16} homologue. The syntheses of most of these compounds are described by Ohloff (supra). As shown in Figure 1, 17 β -hydroxy-5 α -Androstan-3-one (I) and methyl chloroformate (a) in pyridine gives the methyl carbonate, 17 β -methoxycarbonyloxy-5 α -Androstan-3-one (II) which provides a starting material for the 5 α -Androst-16-en-(3-one and 3-ols) (Ohloff, supra at pg 200).

-31-

Alkoxy derivatives are prepared from their corresponding hydroxy steroids by reaction with an alkylating agent such as trimethyloxonium fluoroborate, triethyloxonium fluoroborate or
5 methylfluorosulfonate in an inert chlorocarbon solvent such as methylene chloride. Alternatively, alkylating agents such as alkyl halides, alkyl tosylates, alkyl mesylates and dialkylsulfate may be used with a base such as NaH, KM or KOBut, silver
10 oxide or barium oxide in polar, aprotic solvents as for example, DMF, DMSO and hexamethylphosphoramide.

General procedures for synthetic reactions of steroids are known to those skilled in art. Where time and temperature of reactions must be determined,
15 these can be determined by a routine methodology. After addition of the required reagents, the mixture is stirred under an inert atmosphere and aliquots are removed at hourly intervals. The aliquots are analyzed by chromatography to monitor the
20 disappearance of starting material, at which point the work-up procedure is initiated. If the starting material is not consumed within twenty-four hours, the mixture is heated to reflux and hourly aliquots are analyzed, as before, until the starting material
25 disappears. In this case the mixture is allowed to cool before the work-up procedure is initiated.

Purification of the products is accomplished by means of chromatography and/or crystallization, as known to those skilled in the
30 art.

2. Preparation of 19-OH derivatives

Synthesis of 19-OH-Androsta-4,16-diene-3-one.

-32-

This compound has been disclosed as an intermediate in the synthesis of 19-oxo-3-aza-A-homo-5B-androstane (Habermehl, et al., Z. Naturforsch. (1970) 25b:191-195). A method of synthesizing this
5 compound is provided.

D. Pharmaceutical Compositions and Methods of Use

An embodiment of the subject invention is a method of altering the hypothalamic function of an
10 individual. Another embodiment is altering an autonomic function of an individual. These autonomic functions include but are not limited to heart rate, respiratory rate, brain wave patterns (percentage alpha cortical activity), body temperature. Other
15 embodiments include, but are not limited to, methods of diminishing negative affect, negative mood or negative character traits of an individual. Another embodiment is a method of treating female premenstrual stress. All of these embodiments are
20 accomplished by means of the non-systemic, nasal administration of certain 16-Androstene steroids, combinations of 16-Androstene steroids and combinations of one or more 16-Androstene steroids and one or more Estrene steroids.

25 This particular mode of administration is distinguished from alternative modes, such as ingestion or injection, in several important ways, these by virtue of the direct contact with the VNO provided by the nasal administration of the steroid
30 ligand. In the methods of this invention, the appropriate ligand is administered directly to the chemoreceptors in the nasal passage and the vomeronasal organ, without pills or needles - i.e., non-invasively. Drug action is mediated through

-33-

binding of the ligands, described herein, to specific receptors displayed by neuroepithelial cells in the nose, preferably in the VNO. This Furthermore, the mode of drug action is through the nervous system and not through the circulatory system - thus brain function can be affected without consideration of the blood-brain barrier. These methods of treatment provide a direct means of affecting the hypothalamus through the nervous system because there is only one synaptic junction between pheromone receptors and the hypothalamus. Because sensory nerves are addressed to a specific location in the brain, this method has a highly specific drug effect, thereby greatly reducing the potential of undesirable side-effects.

VNO contact is important because the VNO is associated with chemoreceptive/pheromonal function. The VNO consists of a pair of blind tubular diverticula which are found at the inferior margin of the nasal septum. The VNO contains neuro-epithelia, the axons of which have direct synapses to the amygdala and from there, to the hypothalamus. The existence of the VNO has been well documented in most terrestrial vertebrates including the human fetus; however, in adult humans it is generally thought to be rudimentary (See Johnson, et al., supra).

The ligand substances described herein, or their sulfated, cypionated, benzoated, propionated, or glucuronated derivatives, may be administered directly, but are preferably administered as compositions. They are prepared in a liquid dosage form such as, for example, liquids, suspensions or the like, preferably in unit dosage forms suitable for single administration of precise dosages. Liquid dosages may be administered as nose drops or as an aerosol. Alternatively, the active compound can be

-34-

prepared as a creme or an ointment composition and applied topically within the nasal cavity. As another alternative, delivery may occur by controlled release of these agents by encapsulation either in bulk or at a microscopic level using synthetic polymers, such as silicone, and natural polymers such as gelatin and cellulose. The release rate can be controlled by proper choice of the polymeric system used to control the diffusion rate (Langer, R.S. and Peppas, N.A., *Biomaterials* 2,201, 1981). Natural polymers, such as gelatin and cellulose slowly dissolve in a matter of minutes to hours while silicone remains intact for a period of months. The compositions will include a conventional pharmaceutical carrier or excipient, one or more of the active 16-Androstene compound(s) of Formula I, and the composition may or may not additionally include one or more Estrene steroids. In addition, the compositions may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, etc.

The most likely means of communication of a semiochemical ligand is the inhalation of a naturally occurring pheromone present on the skin of another. Several 16-Androstene steroids, including 5 α -Androst-16-en-3 α -ol and 5 α -Androst-16-en-3-one, 4, 16-Androstadien-3-one, 5 α -Androstadien-3 β -ol, and perhaps 5 α -Androstadien-3 α -ol, are naturally occurring in humans and may be present on the skin. It is estimated that the naturally occurring maximum concentration of a 16-Androstene steroid on human skin is from 2 to 7 ng/cm². During intimate contact it is estimated that a human would be exposed to no more than 700 ng of a naturally occurring steroid. Since these compounds are relatively nonvolatile, it is estimated that, even during intimate contact, a

-35-

human subject would inhale no more than 0.7 pg of a naturally occurring steroid from the skin of another. From the amount inhaled only about 1% would reach the receptors of the vomeronasal organ. Thus the
5 estimated maximum natural exposure to naturally produced pheromones would be 0.007 pg.

The amount of semiochemical ligand administered will of course, be dependent on the subject being treated, the severity of the
10 affliction, the manner of administration, the frequency of administration, and the judgment of the prescribing physician. However, a single dosage of at least about 10 picograms, delivered directly into the lumen of the vomeronasal organ, is effective in
15 eliciting a transient autonomic response. When administered to the nasal cavity, the dosage is about 100 picograms to about 100 micrograms, preferably about 1 nanogram to about 10 micrograms, more preferably about 10 nanograms to 1 about microgram.
20 The frequency of administration is desirably in the range of an hourly dose to a monthly dose, preferably from 8 times/day to once every other day, more preferably 1 to 3 times per day. Ointments containing one or more active compounds and optional
25 pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, can be prepared using a base such as, for example, petroleum jelly, lard, or lanolin.

30 Liquified pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc. an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline,
35 aqueous dextrose, glycerol, ethanol, and the like, to

-36-

thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH
5 buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, etc. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example,
10 see Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, PA, 15th Ed., 1975. The composition or formulation to be administered will, in any event, contain a quantity of one or more of the active compound(s) in an amount effective to
15 alleviate the symptoms of the subject being treated.

For aerosol administration, the active ingredient is preferably supplied in finely divided form along with a surfactant and a propellant. Typical percentages of active ingredients are 0.001
20 to 2% by weight, preferably 0.004 to 0.10%.

Surfactants must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22
25 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, oleostearic and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride such as, for example, ethylene glycol, glycerol, erythritol, arabitol, mannitol,
30 sorbitol, and hexitol anhydrides derived from sorbitol (the sorbitan esters sold under the trademark "Spans") and the polyoxyethylene and polyoxypropylene derivatives of these esters. Mixed esters, such as mixed or natural glycerides, may be
35 employed. The preferred surface-active agents are

-37-

the oleates or sorbitan, e.g., those sold under the trademarks "Arlacel C" (sorbitan sesquioleate), "Span 80" (sorbitan monooleate) and "Span 85" (sorbitan trioleate). The surfactant may constitute 0.1-20% by weight of the composition, preferably 0.25-5%.

The balance of the composition is ordinarily propellant. Liquefied propellants are typically gases at ambient conditions, and are condensed under pressure. Among suitable liquefied propellants are the lower alkanes containing up to five carbons, such as butane and propane; fluorinated or fluorochlorinated alkanes, such as are sold under the trademark "Freon". Mixtures of the above may also be employed.

In producing the aerosol, a container equipped with a suitable valve is filled with the appropriate propellant, containing the finely divided active ingredient and surfactant. The ingredients are thus maintained at an elevated pressure until released by action of the valve.

Yet another means of administration is topical application of a volatile liquid composition to the skin, preferably facial skin, of an individual. The composition will usually contain an alcohol such as ethanol or isopropanol. A pleasant odorant may also be included in the composition.

F. Measuring Affect, Mood and Character Trait.

Feeling states associated with affects, moods and character traits are generally measured by use of a questionnaire. For example questionnaires comprising a number of adjectives which refer to feeling states may be administered to an individual. The individual evaluates his or her feeling state

-38-

described by the adjective and rates the intensity of the feeling on a numerical scale. Clustering of related adjectives and statistical analysis of a subject's evaluation of each adjective provides a basis for the measurement of various feeling states.

Alternatively, feeling states may be measured by autonomic changes, such as those used in polygraphic evaluations (galvanic skin response, pulse rate and the like). Cabanac, M. Annual Review of Physiology (1975) 37:415; Hardy, J.D., "Body Temperature Regulation", Chapter 59, pp. 1417. In: Medical Physiology. Vol. IIEd.: VB Mountcastle (1980); Wolfram Bouscein. Electrodermal Activity (Plenum Press 1992). In addition, non-verbal cues such as facial expression and body posture may be evaluated.

G. Uses in the Treatment of Certain Types of Psychiatric Disorders.

Compositions suitable for topical and/or intranasal administration, and which contain one or more of the steroidal substances described herein, are useful as therapeutics with efficacy in the treatment of certain types of psychiatric disorders, particularly certain types of neuroses. In a preferred embodiment, and unlike other drug formulations designed to invoke a systemic effect, the composition is formulated to minimize the likelihood of mucosal or transdermal absorption since the mode of action of these substances as therapeutics is by stimulation of neural receptors in the nose, more particularly in the VNO. Since the compositions of the instant invention are effective as the result of stimulation of chemosensory receptors in the VNO and not as a result of systemic circulation, absorption and the attendant

-39-

incorporation of the active ingredient or ingredients into systemic circulation would not be efficacious, but rather might actually increase the likelihood of side effects. These compositions are useful when
5 administered in a method such that one or more of the Estrenes or 16-Androstenes of the composition are provided to the VNO of the subject.

Treatable neuroses may be acute and/or transient or they may be persistent or recurrent.
10 They generally involve abnormal symptoms that may include mood changes (anxiety, panic, depression) or limited abnormalities of thought (obsessions, irrational fears) or of behavior (rituals or compulsions, pseudoneurological or hysterical
15 conversion signs). In such disorders, these compositions may have some beneficial effects for short periods, particularly by modifying associated anxiety or depression or the like.

Other characterological disorders may also
20 be treatable by the intranasal administration of the compositions of this invention. These conditions include characteristic personality styles, e.g., paranoid, withdrawn, psychopathic, hypochondrial and the like; or behavioral patterns, e.g., abuse of
25 alcohol or other substances, socially deviant or perverse behavior and the like that may run counter to societal expectations.

Certain physical or physiological conditions, both normal (such as menstruation) and
30 disease or injury related (such as chronic illness) may have a psychological component that manifests as a psychiatric disorder, neurotic disorder, or as an anxious or depressed affect or mood. These conditions are also treatable by methods of
35 administration of the compositions of the present

-40-

invention such that one or more of the estrenes or androstenes of the composition are provided to the VNO of the subject.

i. Anxiety

5 Anxiety is a disagreeable emotional state characterized by feelings of uneasiness, impending danger, apprehension or tension. Compositions suitable for intranasal administration, and which contain one or more of the Estrene steroid substances
10 described herein, are useful as therapeutics with efficacy in the reduction of anxiety. There are numerous manifestations of this disorder ranging from mild perturbations to syndromes which can be incapacitating. Anxiety is not only a cardinal
15 symptom of many psychiatric disorders but also a component of many normal physiological and social conditions. In addition, symptoms of anxiety are commonly associated with depression and especially with dysthymic disorder (neurotic depression) and
20 many personality disorders. Anxiety disorders can be separated and include panic disorders, phobias and generalized anxiety disorders, post-traumatic stress disorder and obsessive-compulsive disorder.

ii. Drugs used in the treatment of anxiety

25 Drug treatment is the major treatment method of anxiety, often used in conjunction with behavior therapy. The drugs used most frequently to treat anxiety are the benzodiazepines. Benzodiazepines have hypnotic, sedative, anxiolytic,
30 anticonvulsant and muscle relaxant actions. Consequently, they are used for many indications and in many different conditions than anxiety disorders (e.g., insomnia, alcohol withdrawal states, muscle spasms, epilepsy, anesthesia and sedation for
35 endoscopic procedures). These drugs are relatively

-41-

safe, and have advantages in comparison with the previously used barbiturates. They have a rapid onset of action, and produces a sense of euphoria. However, despite their good safety record,

5 benzodiazepines are associated with a number of properties that limit their use including sedation, ataxia, memory impairment, reduced motor coordination and dangerous addictive effects, especially when used with alcohol. Both physicians and patients are

10 increasingly aware of this potential for addiction; and there is an evident desire, if not trend, to move away from their longterm use.

Recently, a new class of non-benzodiazepine anxiolytic agents, azaspirodecanediones, which may

15 have more selective anxiolytic properties, has been approved. The first marketed member of this class is buspirone. This compound has been reported to be as effective as benzodiazepines in the treatment of Generalized Anxiety Disorder. Compared to

20 benzodiazepines, buspirone is less sedating, potentiates the effect of ethanol to a lesser degree, and has a low addictive potential. However, in normal clinical use, it has been found to be difficult to substitute buspirone for benzodiazepines

25 in patients already receiving benzodiazepines since buspirone does not create a feeling of euphoria, and patients do not feel as satisfied as when taking benzodiazepines. In addition, buspirone's slow onset of action (1-3 weeks) limits its effectiveness in

30 treating acute anxiety. Systemically administered buspirone anxiolytic has been shown to have a high affinity for the CNS serotonin 5-HT receptor and this is thought to be the mode of anxiolytic action.

Another class of new anxiolytic is the

35 systemically administered benzofuran derivatives.

These substances demonstrate a high affinity for the CNS dopamine D2 receptor.

iii. Mood (Affective) disorders

The compositions of the instant invention
5 are also useful in the treatment of some of the
symptoms of certain types of mood disorders. Mood
disorders such as major depression and mania are
characterized by changes in mood as the primary
clinical manifestation. Either extreme of mood may
10 be associated with psychosis, characterized by
disordered delusional thinking and perceptions, often
congruent with the predominant mood. Conversely,
psychotic disorders may have associated or secondary
changes in mood. Similarly, normal grief, sadness
15 and disappointment and the dysphoria or
demoralization often associated with medical illness
or with normal cyclic dysfunctions such as
premenstrual stress may also be mitigated by the
intranasal application of the compositions of the
20 instant invention.

iv. The mode of action of the compositions
of the instant invention

The mode of action of the compositions of
the instant invention is quite different from any
25 modes of treatment that have been heretofore
reported. The active steroid and steroid-like
compounds described in the instant invention
stimulate and/or bind to receptors in the VNO that
are directly available to topical administration.
30 Direct access to CNS receptors is not required.
Stimulation of these receptors then produces a signal
transmitted through a neural pathway and inducing a

-43-

neuropsychic response, likely in the hypothalamic region of the subject's brain. This response manifests as discrete changes in a variety of autonomic functions, including but not limited to pulse rate, respiratory frequency, E.E.G. patterns, evoked potential, skin temperature, galvanic skin response and pupillary diameter. The response also manifests as a measurable reduction of negative affect and negative mood.

While the relationship between the modification of hypothalamic function and the treatment of psychiatric disorders is not fully understood, it is clear that the compositions of this invention are effective in the reduction of self-evaluative negativity - negative mood characteristics as well as negative character. These changes are accompanied by autonomic changes, effected by the hypothalamus, that are consistent with an increase in parasympathetic tone (or alternatively a decrease in sympathetic tone).

Since the active compounds of this invention are pheromone-like in their specificity, the compounds exhibit a species specificity in their stimulation and activity add therefore demonstration of efficacy cannot be done in animal models.

III. Examples

The following examples are intended to illustrate but not to limit the invention.

Abbreviations used in the examples are as follows: aq.= aqueous; RT.=room temperature; PE=petroleum ether (b.p. 50-70°); DMF=N,N-dimethylformamide; DMSO=dimethyl sulfoxide; THF=tetrahydrofuran.

Example 1 - Androsta-4,16-dien-3-one (4).

10 This synthesis is depicted in Figure 1. Several methods are known for the conversion of testosterone into Androsta-4,16-dien-3-one (Brooksbank et al., Biochem. J. (1950) AZ:36). Alternatively, thermolysis (460°) of the methyl
15 carbonate of testosterone gives Androsta-4,16-dien-3-one in 90% yield. 17B-MethoxyCarbonyloxy-androst-4-en-3-one (IV) was prepared from testosterone (III. Fluka) with methyl chloroformate/pyridine (a) in 76% yield (after recrystallization from MeOH). M.p. 140-
20 141°, $[\alpha]_D = +95.4^\circ$ (c = 1.10) - IR. (CDCl₃): 1740s, 1665s, 1450s, 1280s, - ¹H-NMR. (360 MHz): 0.87 (s, 3 H); 1.20 (s, 3 H); 3.77 (s, 3 H); 4.53 (br. t, J 8, 1 H); 5.75 (s, 1 H). A solution of the methyl carbonate IV in toluene was pyrolyzed (b) as
25 described for I. Recrystallization of the crude

-45-

product from acetone at RT. gave pure ketone 4 in 90% yield. M.p. 127-129.5°, $(\alpha)_D +118.9^\circ$ ($C = 1.32$) ($[\lambda]$: m.p. 131.5-133.5° (hexane), $[\alpha]_D^{16} = +123 \pm 3.5^\circ$ ($C = 1.03$)). - IR. ($CDCl_3$): 3050w, 1660s, 1615m. - 1H -
5 NMR. (360 MHz): 0.82 (s, 3 H); 1.22 (s, 3 H); 5.70 (m, 1 H); 5.73 (s, 1 H); 5.84 (m, 1 H).

Example 2 - Androsta-4,16-dien-3 α -ol (5) and -3 β -ol (6).

These syntheses are depicted in Figure 1.

10 Androsta-4,16-dien-3-one (4) was reduced at -55° with lithium tris (1, 2-dimethylpropyl) hydridoborate in THF (c) as described for the preparation of 2 (Figure 1). Chromatography on silica gel with CH_2Cl_2 /ethyl acetate 9:1 gave pure axial alcohol 5 (48% yield) and
15 pure equatorial alcohol 6 (48% yield). Analytical samples were further purified by recrystallization (from PE at -30° for 5, from cyclohexane at RT. for 6).

Data of 5. M.p. 77-79°, $[\alpha]_D +120.6^\circ$ ($C =$
20 1.26) - IR. ($CDCl_3$): 3620m, 3440m br., 1660m, 1595w. 1H -NMR. (360 MHz): 0.79 (s, 3 H); 1.02 (s, 3 H); 4.07 (m, $w_{1/2} \approx 10$, 1 H); 5.48 (d x d, J 5 and 2, 1 H); 5.71 (m, 1 H); 5.85 (m, 1 H).

Data of 6. M.p. 116.1190, $[\alpha]_D +53.9^\circ$ (C
25 = 1.28) ($[\lambda]$: m.p. 116.1180, $[\alpha]_D^{16} +59.3^\circ$ ($C = 0.4$) -

-46-

IR. (CDCl₃): 3610m, 3420m br., 3050m, 1660m, 1590w.
- ¹H- NMR. (360 MHz): 0.78 (s, 3 H); 1.08 (s, 3 H);
4.15 (m, w_{1/2} ≈ 20, 1 H); 5.30 (m, w_{1/2} ≈ 5, 1 H); 5.71
(m, 1 H); 5.85 (m, 1 H).

5 Example 3 - Androsta-5,16-dien-3 α -ol (7).

This synthesis is depicted in Figure 2. To
a solution of alcohol 8 (545 mg, 2.0 mmol) in acetone
(100 ml) at 0°C under N₂ was added rapidly Jones
reagent (i, 1.5 ml, ca. 4 mmol). After 5 min., the
10 mixture was poured into a dilute phosphate buffer (pH
7.2, 1200 ml) and extracted with ether. The extracts
were washed with sat. aq. NaCl solution, dried
(Na₂SO₄) and evaporated to give mainly Androsta-5,16-
dien-3-one as an oil (567 mg). The crude product was
15 dissolved in THF (7 ml) and reduced with lithium tris
(1,2-dimethylpropyl) hydridoborate (c) at .55° as
described for the preparation of 2. The crude
product (530 mg) was chromatographed on silica gel
(100 g) with CH₂Cl₂/ethyl acetate 4:1 to give 280 mg
20 (51%) of pure α -alcohol 7 (eluted first) and 13 mg of
starting alcohol 8. A small sample of 7 was
recrystallized from acetone/water at RT. M.p. 1380,
[⁸]D -77.5° (c=1.2. - IR. (CDCl₃): 3580m, 3430m,
1665w, 1590w, - ¹H-NMR. (360 MHz): 0.80 (s, 3 H);

-47-

1.06 (s, 3 H); 4.02 (m, $w_{1/2} \approx 8$, 1 H); 5.44 (m, 1 H);
5.72 (m, 1 H); 5.86 (m, 1 H).

Example 4 - Androsta-5,16-dien-3B-ol (8).

This compound was prepared in 73% yield by
5 a known procedure (Marx, A.F., et al., Ger. Offen.
2,631,915; Chem. Abst. 87:23614p (1977)) from
commercial (Fluka) 3B-hydroxy-androst-5-en-17-one
(VII). M.p. 137°, $[\alpha]_D = -71.9^\circ$ ($c=1.5$) ([48]: m.p.
140-141°, $[\alpha]_D = 68^\circ$. - IR. (CDCl_3): 3600m, 3420m
10 br., 1670w, 1590w, - $^1\text{H-NMR}$. (360 MHz): 0.80 (s, 3
H); 1.05 (s, 3 H); 3.53 (m, $w_{1/2} \approx 22$, 1 H); 5.38 (m, 1
H); 5.72 (m, 1 H); 5.86 (m, 1 H). This synthesis is
depicted in Figure 4.

Example 5 - Alternate synthesis of Androsta-4.16-
15 dien-3-one (25).

The following method of synthesis is
depicted in Figure 3:

Dehydroepiandrosterone p-Toluenesulfonylhydrazone
(23)

20 Dehydroepiandrosterone (VII) (14.4 g, 50.0
m mole) and p-toluenesulfonylhydrazide (12.75 g, 68.5
m mole) in dry methanol (300 ml) were heated under
reflux for 20 hours. The mixture was transferred to

-48-

a conical flask and allowed to cool. The crystalline product was filtered under suction and washed with methanol (50 ml). Further crops of product were obtained by sequentially evaporating the filtrate to 5 75 ml and 20 ml, and allowing crystallization each time. Total yield was 21.6 g (95%).

Androsta-5, 16-dien-3 β -ol (24)

Dehydroepiandrosterone p-toluenesulfonylhydrazone (23) (22.8g, 50.0 m mole) in 10 dry tetrahydrofuran (1.0 liters) was cooled in a dry ice/isopropanol bath, The-mixture was stirred while n-butyl lithium (125 ml of 1.6 M solution in hexane, 200 m mole) was added. The mixture was allowed to warm to room temperature and was stirred for 24 15 hours. Water (50 ml) was added with cooling in ice. The mixture was poured into saturated ammonium chloride solution/ice (500 ml) and extracted with ether (x2). The organic layers were washed with saturated sodium bicarbonate solution (500 ml) and 20 saturated sodium chloride solution (500 ml), dried (MgSO₄) and evaporated in vacuo to give the crude product. This was purified by flash chromatography on 190 g silica gel 60, 230-400 mesh, eluting with ethyl acetate/hexane (20:80-50:50) to give

-49-

crystalline material. The product was recrystallized from methanol (45 ml)/3% hydrogen peroxide -(8 ml) washing with methanol (30 ml)/water (8 ml) to give pure product (6.75 g, 50%).

5 Androsta-4, 16-dien-3-one (25)

A solution of 10 g of Androsta-5,16-dien-3 β -ol (24) in 475 cc of toluene and 75 cc of cyclohexanone was distilled (ca. 50 cc of distillate was collected) to eliminate moisture, 5 g of Al(OPrⁱ)₃ in 50 cc of toluene was added and the solution was refluxed for 1 hour. Water then was added, volatile components were removed by steam distillation and the residue was extracted with chloroform. Evaporation of the dried extract, followed by crystallization of the residue from chloroform-hexane, yielded 7.53 g of Androsta-4,16-dien-3-one (25). Another 0.97 g (total, 8.5 g, 86%) was obtained by chromatography of the mother liquor on neutral alumina.

Example 6 - Synthesis of Androsta-3,5,16-trien-3-yl methyl ether (12).

To a partial solution of androsta-4,16-dien-3-one (1.00 g, 3.70 mmol) in 2.2-dimethoxypropane (5.0 mL, 41 mmol) and 5 mL DMF were added methanol (0.2 mL) and p-toluenesulfonic acid

-50-

monohydrate (26.4 mg, 0.139 mmol). The mixture was refluxed 5 h, after which it was cooled and sodium bicarbonate (152.5 mg) was added. The suspension was partitioned between 50 mL of ice water and 50 mL of ethyl acetate. The organic layer was washed with two 50 mL portions of water + 50 mL of brine, dried over magnesium sulfate, filtered, and concentrated under reduced pressure. The residual oil was taken up in 50 mL of hot hexane and filtered through a 12 mm x 30 mm column of silica gel 60 using 150 mL of hot hexane. The combined filtrates were concentrated under reduced pressure and recrystallized from acetone/methanol to give white crystals (468.0 mg, 1.645 mmol, 44%), m.p. 83-92°C.

15 Example 7 - Synthesis of 17-methylene-Androst-4-en-ols.

To 20-homoandrosta-4,17-dien-3-one (119.0 mg, 0.4184 mmol) in 5 mL of methanol were added sodium borohydride (6.0 mg, 0.16 mmol) and 77 μ L of water. After stirring 2 h further sodium borohydride (32.0 mg, 0.846 mmol) was added and the mixture was stirred overnight. After concentrating under reduced pressure the residue was purified by preparative TLC (5% ethyl acetate/hexane on silica gel) to give a

-51-

more polar (59.8 mg) and a less polar (1.7 mg) product.

Example 8 - Synthesis of 17-methylene-6-oxo-Androsta-4-en-3-one.

5 To a cooled solution of 20-homoandrosta-5,17-dien-3-ol (399.4 mg, 1.394 mmol) in 50 mL of acetone was added 2.67M Jones reagent (2.0 mL, 5.3 mmol). After stirring 1 h the reaction was quenched with isopropanol (1.0 mL, 13 mmol) and poured into
10 100 mL of water. The mixture was extracted three times with 50 mL portions of ethyl acetate and the combined organic extracts were washed with 50 mL of saturated sodium bicarbonate + 50 mL of brine. The organic phase was then dried over magnesium sulfate,
15 filtered, and concentrated under reduced pressure. The residue was recrystallized from 95% ethanol to give an almost white powder (177.8 mg, 0.5958 mmol, 43%), m.p. 113-115°C.

Example 9 - Synthesis of 6 β -OH-Androsta-4,16-dien-3-one.

20 To a solution of androsta-3,5,16-trien-3-yl methyl ether, (12) (200.5 mg, 0.7049 mmol), in 5 mL of 1,2-dimethoxyethane (DME) and 1 mL of water was added m-chloroperbenzoic acid (MCPBA, 77.4%, 173.2

-52-

mg, 0.776 mmol) suspended in 5 mL of DME + 1 mL of water + 0.40 g of 5% (w/w) NaOH dropwise, with stirring, over a period of 90 min. After stirring 18 h further MCPBA (247.0 mg, 1.11 mmol) suspended in 10 mL of DME + 2 mL of water + 0.8 g of 5% (w/w) NaOH was added dropwise, with stirring, over 1 1/2 h. The reaction mixture was stirred 1/2 h and then poured into 25 mL of saturated sodium bicarbonate. The aqueous mixture was extracted three times with 25 mL of ether and the combined organic extracts were washed with 50 g of 5% (w/w) sodium thiosulfate + three 50 mL portions of brine, dried over magnesium sulfate, filtered through Celite, and concentrated under reduced pressure. The resulting crystalline residue was purified by preparative TLC (35% ethyl acetate/hexane on silica gel) followed by two-fold recrystallization from aqueous ethanol to give lustrous white platelets (102.3 mg, 0.3571 mmol, 51%), m.p. 165-166°C.

20 Example 10 - Electrophysiology of Androstane
Stimulation of the Human VNO and Olfactory
Epithelium.

A non-invasive method has been employed to record local electrical potentials from the human vomeronasal organ (VNO) and from the olfactory

25

-53-

epithelium (OE). Localized gaseous stimulation was applied to both nasal structures at different instances using specially designed catheter/electrodes connected to a multichannel drug delivery system. This electrode and delivery system has been described by Monti and Grosser (J. Steroid Biochem. and Molec. Biol. (1991) 39:573) and in commonly owned, copending USSN 07/771,414, incorporated herein by reference. The local response of the VNO and the OE showed a correlation with the concentration of the ligand stimulus.

The study was performed on ten clinically normal (screened) volunteers - 2 males and 8 females, ranging in age from 18 to 85 years. The studies were conducted without general or local anesthetics.

The catheter/electrodes were designed to deliver a localized stimulus and simultaneously record the response. In the case of VNO recording, the right nasal fosa of the subject was explored using a nasoscope (nasal specule) and the vomeronasal opening was localized close to the intersection of the anterior edge of the vomer and the nasal floor. The catheter/electrode was gently driven through the VNO-opening and the electrode tip placed in the organ's lumen at 1 to 3 mm from the opening. The nasoscope was then removed. In the case of the OE,

-54-

recording the procedure was similar except the positioning of the catheter/electrode was gently placed deep in the lateral part of the medial nasal duct, reaching the olfactory mucosa.

5 Localized gaseous stimulation was done through the catheter/electrode. A constant stream of clean, nonodorous, humidified air at room temperature was continuously passed through a channel of the stimulating system. The stimulating ligand
10 substances were diluted in propylene glycol, mixed with the humidified air, and puffed for from 1 to 2 seconds through the catheter/electrode. It is estimated that this administration provides about 25 pg of the steroid-ligand to the nasal cavity.

15 The results of this study are presented in Figures 4A, 4B, and 4C. The response is measured in millivolt-seconds (mV x s). Androsta-4,16-dien-3-one elicits a significantly stronger VNO response in females than do the other compounds tested (Fig. 4A).
20 Furthermore, the VNO response to Androsta-4,16-dien-3-one is sexually dimorphic - twice as strong in females as it is in males (Fig. 4B). In contrast, the OE response in both males and females is low compared to a strong odorant such as clove (Fig. 4C).

-55-

Example 11 - Measurement of the Change in Receptor Potential of the Neuroepithelium of the VNO in Response to Various Steroids.

The change in receptor potential in response to five different ligands was measured in 40 female (Fig. 5A) and 40 male (Fig. 5B) subjects. Each subject was administered 60 pg of each of seven substances as indicated in the figure. The substances were administered separately for 1 second, using the procedure described in Example 10. The change in potential of the neuroepithelium of the VNO was recorded over time and the integral of the change in potential for each of the forty subjects was averaged. The results are shown in the figure.

Comparison of Figures 5A and 5B show that each steroid is sexually dimorphic in its activity, and that some ligand substances are stronger in males while others are stronger in females.

Example 12 - Measurement of Autonomic Responses to 16-Androstene Stimulation of the VNO.

Various autonomic parameters were monitored as Androsta-4,16-dien-3-one was administered to 40 female subjects using the procedure described in Example 10. Propylene glycol was also administered as a control. The ligand was administered as a 1

-56-

second pulse. The change in autonomic function was first noted within 2 seconds and lasted for up to 45 seconds. As shown in Figure 6, when compared to a propylene glycol control, the Androstane induced a significant change in the integrated receptor potential in the VNO (6A), galvanic skin response (6B), skin temperature (6C), the percentage of cortical alpha wave activity as measured by electroencephalogram (6D), peripheral arterial pulse (6E), and respiratory frequency (6F).

Example 13 - Comparison of the Change in Receptor Potential Induced by Two Androstane Steroids.

60 picograms of each ligand steroid and of a propylene glycol control were administered to 5 female subjects as described in Example 10. As shown in Figure 7, Androsta-4,16-dien-3 β -ol induced a greater change in receptor potential than did Androsta-4,16-dien-3-one.

Example 14 - Psychophysiological Effect of Androstane Stimulation of the VNO.

The psychophysiological effect of Androstane stimulation of the VNO was measured by the coordinate administration of pheromone and questionnaire evaluation of the subject before and

-57-

after administration. The questionnaire included a panel of adjectives used as part of the standard Derogatis Sexual Inventory evaluation.

The subjects were 40 women between the ages of 20 and 45, all in good health. The women were randomly assigned - 20 exposed to placebo and 20 exposed to about 20 picograms of Androsta-4,16-dien-3-one, administered as described in Example 10, supra. Subjects were given a 70 item questionnaire evaluating feeling states immediately before and 30 minutes after administration of either placebo or experimental substance. The 70 adjectives of the questionnaire were randomly administered and subsequently clustered for evaluation based on their relevance to each mood, feeling, or character trait. The results were as follows: Changes in feelings of social warmth, personal well-being, arousal/excitement, and aggression, from before administration to 30 minutes after administration, were not significant in those exposed to the 16-Androstene compared to the changes resulting from administration of the control. However, the decrease in negative affect (nervous, tense, ashamed, anxious, irritable, angry, enraged - T-test: $p < 0.0001$, Anova: $p < 0.04$), negative mood and character (sensitive, regretful, blameworthy, guilty, remorseful, sad,

-58-

hopeless, resentful, worthless, miserable, unhappy, bitter, timid - T-test: $p < 0.0004$, Anova: $p < 0.06$), and overall negativity (the combination of affect and character - T-test: $p < 0.0003$, Anova: $p < 0.05$) were highly significant after 16-Androstene administration as compared to administration of the control.

Overall, these results suggest a sedative and/or anti-anxiety, and/or anti-depressant effect of Androsta-4, 16-dien-3-one when administered intranasally.

Example 15 - Treatment of Women for Premenstrual Stress.

Women experiencing the symptoms of premenstrual stress (PMS) are provided with a pharmaceutical preparation of an Androstane steroid (preferably Androsta-4,16-dien-3-one, or Androsta-4,16-dien-3 α (β)-ol) suitable for nasal administration. The steroid is provided as an ointment at a concentration of about 1 microgram/ml and about 0.1 ml is applied. The ointment is applied just inside the nare of each nostril, three times daily. A similar method of treating PMS uses an aerosol preparation of the same steroid. The aerosol is sprayed into each nostril three times daily.

Example 16 - Electrophysiological Studies

The following electrophysiological studies were performed in 60 clinically normal human volunteers of both sexes (30 male and 30 female) whose ages ranged from 20 to 45 years. No anesthetics were used, and female subjects were excluded if pregnant.

The stimulation and recording system consists of a "multifunctional miniprobe" described elsewhere (Monti-Bloch, L. and Grosser, B.I. (1991) "Effect of putative pheromones on the electrical activity of the human vomeronasal organ and olfactory epithelium," J. Steroid Biochem. Molec. Biol. 39:573-582.). The recording electrode is a 0.3 mm silver ball attached to a small (0.1 mm) silver wire insulated with Teflon®. The surface of the electrode is first treated to produce a silver chloride interface, and is then covered with gelatin. It is positioned within a small caliber Teflon® catheter (dia = 5 mm) such that the tip of the electrode protrudes approximately 2 mm. The Teflon® catheter is 10 cm in length and constitutes the terminal extension for a multichannel delivery system which delivers a continuous air stream carrying discreet pulses of chemosensory stimuli. The air stream first passes into a small chamber and is bubbled through a

-60-

solution containing either a vomeropherin or an olfactant in a diluent or the diluent alone. A solenoid is used to rapidly redirect the air stream from the chamber to a route which bypasses the chamber. This creates a discreet pulse of stimulant in the air stream. A second, outer Teflon® tube with a diameter of 2 mm surrounds the catheter-electrode assemblage, and its central end is connected to an aspirator that provides continuous suction of 3ml/s. This concentric arrangement of the outer suction tube allows the emitted chemosensory stimuli to be localized to an area we call a "minifield" (approx. dia = 1 mm), and it avoids diffusion of substances either to the area outside the intended stimulation site or into the respiratory system. The entire stimulating and recording assemblage may be positioned either on the neurosensory epithelium within the VNO, or on the surface of the olfactory or respiratory epithelium.

20 Electro-vomeronasogram (EVG): Recordings are carried out in a quiet room with the subject supine; the multi-functional miniprobe is initially stabilized within the nasal cavity using a nasal retractor placed in the vestibule. Reference and ground electrodes consist of silver discs (8 mm), both of which are positioned on the glabella.

-61-

The entrance to the VNO, or vomeronasal pit, is identified by first dilating the nasal aperture and vestibule. A 6x magnifying binocular loupe with halogen illumination is then used to
5 introduce the tip of the Teflon® catheter and recording electrode assemblage into the VNO opening where it is stabilized at an approximate depth of 1 mm within the vomeronasal passage. Optimal placement of the recording electrode is signaled after testing
10 for an adequate depolarization in response to a test substance.

Electrical signals from the recording electrode are fed to a DC amplifier after which they are digitized, computer monitored, and stored. The
15 peak-to-peak amplitude of the signals is measured, and the area under the depolarization wave is integrated, while continuously monitoring the signal both on the computer screen and on a digital oscilloscope. Artifacts produced by respiratory
20 movements are deleted by training the subjects to practice mouth breathing with velopharyngeal closure.

Chemosensory Stimulants: Olfactory test substances are cineole, and 1-carvone; vomeropherins are A, B, C, D, E and F. Samples of vomeropherins in
25 concentration of 25-800 fmoles are delivered in the continuous air stream for durations from 300

-62-

milliseconds to 1 second. Usually, intervals of 3 to 5 minutes separated each series of short test pulses. All components of the lines carrying the test stimuli are made of Teflon®, glass or stainless steel and are carefully cleaned and sterilized before each use.

Electro-olfactogram (EOG): Olfactory recordings employed the same stimulating and recording multifunctional miniprobe as that used for the VNO. The tip was slowly introduced until the recording electrode touched the olfactory mucosa. Adequate placement was signaled by a depolarization in response to a pulse of the odorant test substance.

Cortical evoked activity was induced by VNO stimulation with vomeropherins, and olfactory stimulation with odorants delivered in 300 ms air pulses. It was recorded using standard electroencephalographic (EEG) electrodes placed at positions Cz-A1 and Tz-A1 of the international 10120 system; the ground electrode was placed on the mastoid process. Electrodermal activity (EDA) was recorded using standard 8 mm silver electrodes in contact with palmar skin of the medial and ring fingers respectively, through a conductive gel interface. Skin temperature (ST) was recorded by a small (1.0 mm) thermistor probe placed in the right ear lobe. Peripheral arterial pulse (PAP) was

-63-

monitored with a plethysmograph attached to the tip of the index finger. Respiratory frequency (RF) was measured with an adjustable strain gauge placed around the lower thorax. All electrical signals were
5 DC amplified, digitized (MP-100, Biopac Systems) and continuously monitored utilizing a computer.

Statistical Analysis: EVGs or EOGs, peak-to-peak changes and frequency changes of other parameters were measured and statistically analyzed. The
10 significance of the results was determined by either using paired t-tests or analysis of variance (ANOVA).

Effect of Vomeroepherins on the EVG: Each of the vomeroepherins was found to produce a sexually dimorphic receptor potential (Fig. 8A-B). Recordings
15 of the EVG were performed on 30 men and 30 women (ages 20 to 45). Vomeroepherins were diluted and applied as 1 second pulses to the VNO with b minute intervals between pulses when questioned, the subjects were not able to "smell" or otherwise
20 consciously detect any of the vomeroepherins. This finding is in agreement with results previously reported (Monti-Bloch, L. and Grosser, B.I. (1991) "Effect of putative pheromones on the electrical activity of the human vomeronasal organ and olfactory
25 epithelium," J. Steroid Biochem. Molec. Biol. 39:573-582.) which indicated that neither olfactory nor

-64-

vomeropherin test stimuli delivered to the VNO elicit a perceptible sensation at the delivered concentration.

Fig. 8A shows the average response of male subjects (ages 20 to 38) to the diluent, and to equimolar quantities (100 fmoles) of five vomeropherins (A, B, C, D and F), and to E, a stereoisomer of F. The profile of the response to each of the substances was similar in all subjects regardless of age, and no significant differences were revealed either by t-tests or by analysis of variance. For example, A, C and D produced significant effects ($M_{15} = 11.4$ mV, $SD = 3.6$ mV; $M_{76} = 6.4$ mV, $SD = 2.5$ mV, and $M_{84} = 15.1$ mV, $SD = 4.9$ mV; $p < 0.01$), that were consistent in all individual cases. Other vomeropherins depolarized the VNO-receptors to a much lesser extent, but with consistent mean response amplitudes from individual to individual. Vomeropherins active in male subjects produced larger responses than the diluent ($p < 0.001$). B, F and similar concentrations of olfactants induced significantly reduced responses in the male VNO (Fig. 8A and Fig. 9).

A similar experimental protocol was followed with the 30 female subjects (ages 20 - 45). Among the vomeropherins, F (100 fmoles) produced the

-65-

most significant differences within the group (Fig. 8B). Here, A induced a small effect that was significantly different from F ($p < 0.01$). In both populations of subjects, active vomeropherins induced
5 receptor responses having large standard deviations (Fig. 8). When the frequency distribution of the effects of A and F was studied in males and females respectively, we found a bimodal distribution. The significance of this observation is being studied at
10 this point.

E, a stereoisomer of F, does not stimulate the VNO in female subjects while F does (Fig. 8B). This is a demonstration of the specificity of VNO recognition of vomeropherins. In this regard it is
15 interesting to note that while F is a superior vomeropherin, E generates a stronger olfactory effect than does F (Fig. 8B and Fig. 9).

Effects of Vomeropherins on the EOG: The summated receptor potential from the olfactory epithelium (OE)
20 was recorded in 20 subjects: 10 males and 10 females. In contrast to the sensitivity of the VNO to vomeropherins, the OE is less sensitive to these substances. This is true for both males and females (Fig. 9A). The mean receptor potential amplitude
25 ranged from 2.3 mV to 0.78 mV. In this study, B was the only vomeropherin having significant effect in

-66-

the OE ($p < 0.02$). Of the subjects questioned about odorant sensations following each stimulus presentation, 16 reported no olfactory sensation, while three males and one female described B as an unpleasant odor. This finding reveals that at the concentrations used in our study, most vomeropherins are not effective stimulants of the olfactory receptors, but do have a clear effect on vomeronasal receptors.

- 10 Effects of Olfactants on the EVG and EOG: In contrast to vomeropherins, the olfactants 1-carvone and cineole produce only a minor local response in the VNO (Fig. 9B). This was true for both men and women. As expected, these olfactants produced a strong response in both men and women ($p < 0.01$) when locally applied to the OE (Fig. 9A). The diluent depolarized olfactory receptors to a lesser extent than cineole or 1-carvone ($p < 0.01$), and it did not produce an olfactory sensation.
- 15
- 20 Reflex Effects of Vomeropherins: Studies were conducted to determine the central nervous system (CNS) reflex responses to vomeropherin stimulation of the VNO. The sexually dimorphic local responses induced by vomeropherins (Fig. 8A and B) were mirrored in the autonomic response of male & female subjects. In male subjects (Fig. 8C), A and C
- 25

-67-

decreased skin resistance (electrodermal activity = EDA) ($p < 0.01$, $n = 30$). In female subjects. (Fig. 8B), F and B produced greater decrease in EDA than A or C ($p < 0.01$, $n = 30$).

5 Vomeropherins A and C induced a significant increase in skin temperature (ST) (Fig. 8G) in 30 male subjects ($p < 0.01$); however D induced significant temperature decrease ($p < 0.01$). In 30 female subjects (Fig. 8H) B and F evoked a significant increase in
10 skin temperature (ST) ($p < 0.01$) compared to A and C. In female subjects vomeropherins produced changes in EDA and ST with a greater standard deviation than in males.

Cortical activity was recorded from Cz and
15 Tz in male and female subjects during application to the VNO of air pulses (300 ms to 1 sec) containing 200 fmoles of vomeropherin (Fig. 8G and H). In males (Fig. 8E) A, C and D significantly increased alpha cortical activity with a latency of 270-380 ms. D
20 and A evoked the strongest effect ($p < 0.01$).

Synchronization of the EEG was sustained for 1.5 to 2.7 minutes after application of a single pulse of active substance. In females (Fig. 8F), a single pulse (200 fmoles) of B or F applied to the VNO
25 increased alpha cortical independent of the response of olfactory receptors. We found characteristic

-68-

specificities in the response of the human VNO and the olfactory epithelium which suggests that they are independent functional systems with separate connections to the CNS (Brookover, C. (1914) The
5 nervus terminalis in adult man. J. Comp. Neurol. 24:131-135.) There is also preliminary evidence that the EVG is not associated with trigeminal nociceptor endings since application of a local anesthetic (2% lidocaine) to the respiratory epithelium of the nasal
10 septum neither blocks nor diminishes the EVG (Monti-Bloch, L. and Grosser, B.I. (1991) "Effect of putative pheromones on the electrical activity of the human vomeronasal organ and olfactory epithelium," J. Steroid Biochem. Molec. Biol. 39:573-582.), also,
15 subjects failed to report sensations of pain as a consequence of any of the stimulation procedures.

VNO receptors are clearly more sensitive to vomeropherins than to any of the olfactants tested; the opposite is true for olfactory receptors. While
20 the OE may have receptor sites for some vomeropherins, the response specificity of the VNO is clearly different.

Sexual differences were noted in the specificities and effects of two groups of
25 vomeropherins, A, C and D; and B and F. This suggests a possible receptor-related sexual

-69-

dimorphism. The findings suggest the activation of components of the autonomic nervous system in the adult human by vomeropherin stimulation of the VNO.


Furthermore, the results suggest that
5 stimulation of the VNO with vomeropherins produces
synchronization of the EEG (Fig. 8G and H) Thus, the
evidence herein indicates that the vomeronasal system
responds to a variety of chemosensory stimuli, and
that some are able to induce reflex autonomic
10 activity.

-70-

CLAIMS

We Claim:

1. A pharmaceutical composition suitable for nasal administration in an individual, said composition comprising an Androstane steroid and a pharmaceutically acceptable carrier, wherein said steroid has the formula:



- wherein P₁ is selected from the group consisting of oxo, α -(β -) hydroxy, α -(β -) acetoxy, α -(β -) propionyloxy, α -(β -) methoxy, α -(β -) lower acyloxy, α -(β -) lower alkyloxy, and α -(β -) benzoyloxy; P₂ is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl, alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P₃ is absent or is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl,

-71-

- alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P_4 is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P_5 represents one or 2
- 5 substituents, wherein P_5 comprises one or two hydrogen atoms, methyl, methylene, or one or two halo atoms; P_6 is hydrogen or halo; and "a", "b", "c", "d", "e", "f", and "h" are alternative sites for optional double bonds.
- 10 2. A composition according to Claim 1 wherein "b" is a double bond.
3. A composition according to Claim 2 wherein "c" or "d" is a double bond.
4. A composition according to Claim 1
- 15 wherein "a" and "c" are double bonds.
5. A composition according to Claim 1 wherein P_3 is methyl, "h" is an optional double bond, and P_5 is methylene or one or two hydrogen atoms.
6. A composition according to Claim 1
- 20 wherein P_3 is methyl and "h" is a double bond.

-72-

7. A composition according to Claim 6 wherein said steroid is selected from the group consisting of Androsta-5,16-dien-3 α -ol, Androsta-4,6,16-triene-3-one; Androsta-4,16-dien-3,6-dione; 5 Androsta-4,16-dien-19-ol-3-one; 3-Methoxy-androsta-3,5,16-triene; and Androsta-4-16,-dien-6 α -ol-3-one.

8. A composition according to Claim 1 wherein P₃ is methyl.

9. A composition according to Claim 8
10 wherein said steroid is Androst-4-en-3-one.

10. A composition according to Claim 1 wherein P₃ is methylene.

11. A composition according to Claim 10 wherein said steroid is selected from the group
15 consisting of 17-Methylene-androst-4-en-3 α -ol; 17-Methylene-androst-4-en-3 β -ol; and 17-Methylene-6-oxo-androst-4-en-3-one.

12. A composition according to Claim 1 wherein P₃ is methyl and "f" is a double bond.

-73-

13. A composition according to Claim 12 wherein said steroid is 10 α ,17-Dimethyl-gona-4,17-dien-3-one.

14. A pharmaceutical composition according to Claim 1 wherein "a" or "b" is a double bond.

15. The pharmaceutical composition of Claim 1 wherein said steroid is selected from the group consisting of Androsta-4,16-dien-3-one, Androsta-4,16-dien-3 α -ol and Androsta-4, 16-dien-3 β -ol, or mixtures of two or more thereof.

16. The pharmaceutical composition of any of Claims 1 through 15 wherein said steroid is dissolved in said carrier.

17. The pharmaceutical composition of any of Claims 1 through 15 wherein said composition is in a liquid form.

18. The pharmaceutical composition of any of Claims 1 through 15 wherein said composition further contains a pharmaceutically acceptable ointment base.

-74-

19. The pharmaceutical composition of any of Claims 1 through 15 which contains no more than one of said steroids.

20. The pharmaceutical composition of any of Claims 1 through 15 which contains more than one of said steroids.

21. The pharmaceutical composition of any of Claims 1 through 15 additionally comprising an Estrene steroid.

10 22. A method of altering a hypothalamic function of an individual, said method comprising:

providing a ligand for a chemoreceptor
displayed on the surface of nasal
neuroepithelial cell of said individual
15 wherein said cell is a part of tissue other
than olfactory epithelia; and,

administering said ligand within a nasal
passage of said individual such that said
ligand binds specifically to said receptor
20 and results in an alteration of
hypothalamic function of said individual.

-75-

23. A method of altering an autonomic function of an individual, said method comprising:


5 providing a ligand for a chemoreceptor of a nasal neuroepithelial cell of-said individual wherein said cell is a part of tissue other than olfactory epithelia; and, administering said ligand within a nasal passage of said individual such that said ligand binds specifically to said receptor
10 and results in an alteration of hypothalamic function of said individual.

24. The method of Claim 23 wherein said neuroepithelial cell is located within a vomeronasal organ of said individual.

15 25. The method of claim 24 wherein said ligand comprises an Androstane.

26. The method of claim 25 wherein said Androstane has the formula:

-76-



wherein P_1 is selected from the group consisting of oxo, α -(β -) hydroxy, α -(β -) acetoxy, α -(β -) propionyloxy, α -(β -) methoxy, α -(β -) lower acyloxy, α -(β -) lower alkyloxy, and α -(β -) benzoyloxy; P_2 is

5 selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl, alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P_3 is absent or is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl,

10 alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P_4 is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P_5 represents one or 2

substituents, wherein P_5 comprises one or two

15 hydrogen atoms, methyl, methylene, or one or two halo atoms; P_6 is hydrogen or halo; and "a", "b", "c", "d", "e", "f", and "h" are alternative sites for optional double bonds.

-77-

27. A method according to Claim 26 wherein "b" is a double bond.

28. A method according to Claim 27 wherein "c" or "d" is a double bond.

5 29. A method according to Claim 26 wherein "a" and "c" are double bonds.

30. A method according to Claim 26 wherein P₃ is methyl, "h" is an optional double bond, and P₅ is methylene or one or two hydrogen atoms.

10 31. A method according to Claim 26 wherein P₃ is methyl and "h" is a double bond.

32. A method according to Claim 31 wherein said steroid is selected from the group consisting of Androsta-5,16-dien-3 α -ol, Androsta-4,6,16-triene-3-one; Androsta-4,16-dien-3,6-dione; 19-Hydroxy-
15 androsta-4,16-dien-3-one; 3-Methoxy-androsta-3,5,16-triene; and 6 α -Hydroxy-androsta-4-16,-dien-3-one.

33. A method according to Claim 26 wherein P₃ is methyl.

-78-

34. A method according to Claim 33 wherein said steroid is Androst-4-en-3-one.

35. A method according to Claim 26 wherein P_5 is methylene.

5 36. A method according to Claim 35 wherein said steroid is selected from the group consisting of 20-Homo-androsta-4,17-dien-3 α -ol; 20-Homo androsta-4,17-dien-3 β -ol; and 20-Homo androsta-4,17-dien-3,6-dione.

10 37. A method according to Claim 26 wherein P_5 is methyl and "f" is a double bond.

38. A method according to Claim 37 wherein said steroid is 10 α ,17-Dimethyl-gona-4,17-dien-3-one.

39. The method of Claim 26 wherein "a" or
15 "b" is a double bond.

40. The method of Claim 26 wherein at least one Androstane steroid is selected from the group consisting of Androsta-4, 16-dien-3-one, Androsta-4,16-dien-3 α -ol and Androsta-4,16-dien-3 β -
20 ol.

-79-

41. The method of Claim 40 wherein said steroid is Androsta-4, 16-dien-3-one.

42. The method of any of Claims 22 through 40 wherein the amount of said ligand that is administered is at least about 100 picograms, but no more than about 100 micrograms.

43. The method of Claim 42 wherein the amount of said ligand that is administered is at least about 1 nanograms, but no more than about 10 micrograms.

44. The method of Claim 43 wherein the amount of said ligand that is administered is at least about 10 nanograms, but no more than about 1 microgram.

15 45. The method of any of Claims 22 through 40 further comprising preparing a pharmaceutical composition of said ligand dissolved in a pharmaceutically acceptable carrier.

46. The method of Claim 45 wherein said 20 pharmaceutical composition is an ointment.

-80-

47. The method of Claim 45 wherein said pharmaceutical composition is liquid.

48. The method of Claim 45 wherein the administration is by aerosol.

5 49. The method of any of Claims 22 through 40 wherein more than one Androstane steroid is administered.

 50. The method of any of Claims 22 through 40 further comprising nasally co-administering to
10 said individual an Estrene steroid.


 51. The method of any of Claims 22 through 40 wherein said function is the diminution of negative affect.

 52. The method of any of Claims 22 through
15 40 wherein said individual is a woman.

 53. The method of Claim 52 wherein said alteration of hypothalamic function results in a reduction of premenstrual stress in women.

-81-

54. The method of any of Claims 22 through 40 wherein said function is the alleviation of the symptoms of psychoses, depression or anxiety.









FIG. 3.



*FIG. 4B.*

4/10




FIG. 5A.

5/10










FIG. 5B.

6/10

**FIG. 6A.****FIG. 6B.****FIG. 6C.****FIG. 6D.****FIG. 6E.****FIG. 6F.**

7/10




FIG. 7.




FIG. 9A.




FIG. 9B.

8/10




FIG. 8A.




FIG. 8B.




FIG. 8C.




FIG. 8D.




FIG. 8E.




FIG. 8F.




FIG. 8G.




FIG. 8H.

9/10

FIG. 10A




FIG. 10B.





FIG. 10C.



10/10

FIG. IIA.




FIG. IIB.





FIG. IIC.



PATENT COOPERATION TREATY

UNITED STATES
International Preliminary
Examining Authority
(IPEA/US)

In re)
International Patent)
Application)
No: PCT/US93/09349)
)
Filed: 28 September 1993)
)
Applicant:)
PHERIN CORPORATION)
)
Title: ANDROSTENE STEROIDS)
AS NEUROCHEMICAL)
INITIATORS OF CHANGE IN)
HUMAN HYPOTHALAMIC)
FUNCTION AND RELATED)
PHARMACEUTICAL)
COMPOSITIONS AND METHODS)
)

July 27, 1994


File No: 18136-0042

"EXPRESS MAIL" MAILING LABEL
NUMBER TB231658625US
DATE OF DEPOSIT July 27, 1994

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS
BEING DEPOSITED WITH THE UNITED STATES
POSTAL SERVICE "EXPRESS MAIL POST OFFICE
TO ADDRESSEE" SERVICE UNDER 37 CFR 1.10 ON
THE DATE INDICATED ABOVE AND IS ADDRESSED
TO: THE COMMISSIONER OF PATENTS AND
TRADEMARKS, WASHINGTON, DC 20231.

TYPED NAME JESSICA KANY

SIGNED



REQUEST FOR RECTIFICATION UNDER PCT RULE 91.1(f)

Honorable Commissioner of Patents
and Trademarks
Washington, DC 20231

Sir:

This is in response to the Notification of
Decision Concerning Request for Rectification

regarding the referenced application mailed May 2, 1994.

Applicant hereby requests, pursuant to Rule 91.1(f), that the International Bureau publish the subject Request for Rectification together with the international application.


The Commissioner is authorized to charge the required fee to Deposit Account 08-1641, Order No. 18136-0042.

The Commissioner is authorized to charge any additional fees or credit any overpayment to Deposit Account 08-1641, Order No. 18136-0042.

Respectfully submitted,


HELLER, EHRLMAN, WHITE
& MCAULIFFE

By


Reginald J. Snyat
Registration No. 28,172

333 Bush Street
San Francisco, California 94104-2878
(415) 772-6432

the 13-position and at the 10-position. Androstenes are a subset of Androstanes and have at least one double bond. Ohloff, G. et al. (Helv. Chim. Acta (1983) 66:192-217), which is incorporated herein by
5 reference, have shown that several members of this group of steroids have an odor which varies with different isomeric, diastereomeric, and enantiomeric forms. Some members of this group have been reported to act as a pheromone in some mammalian species - for
10 instance, 5 α -androst-16-en-3-one and 5 α -androst-16-en-3 α -ol in pigs (Melrose, D.R., et al., Br. vet. J. (1971) 127:497-502). These 16-Androstenes produced by the boar induce mating behavior in estrus sows (Claus, et al., Experimentia (1979) 35:1674-1675).
15 Some studies have noted that, in some species, various characteristics of certain 16-Androstenes (including 5 α -Androst-16-en-3 α -2-ol and 5 α -Androst-16-en-3-one), such as concentration, metabolism, and localization, are sexually dimorphic
20 (Brooksbank et al., J. Endocr. (1972) 52: 239-251; Claus, et al., J. Endocr. (1976) 68:483-484; Kwan, et al., Med. Sci. Res. (1987) 15:1443-1444). For instance, 5 α -Androst-16-en-3 α -ol and 5 α -Androst-16-en-3-one, as well as Androsta-4,16-dien-3-one, have
25 been found at different concentrations in the peripheral blood, saliva and axillary secretions of men and of women (Kwan, T.K., et al., Med. Sci. Res. (1987) 15:1443-1444), and their function as a human pheromone, to the extent of affecting choice and
30 judgement, has been suggested (Id.; see also Gower, et al., "The Significance of Odorous Steroids in Axillary Odour", In, Perfumery, pp. 68-72, Van Toller and Dodd, Eds., Chapman and Hall, 1988); Kirk-Smith, D.A., et al., Res. Comm. Psychol. Psychiat. Behav.
35 (1978) 3:379). Androstenol (5 α -androst-16-en-3 α -ol)



wherein P₁ is selected from the group consisting of oxo, α-(β-) hydroxy, α-(β-) acetoxy, α-(β-) propionyloxy, α-(β-) methoxy, α-(β-) lower acyloxy, α-(β-) lower alkyloxy, and α-(β-) benzoyloxy; P₂ is
 5 selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl, alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P₃ is absent or is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl,
 10 alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P₄ is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P₅ represents one or 2 substituents, wherein P₅ comprises one or two
 15 hydrogen atoms, methyl, methylene, or one or two halo atoms; P₆ is hydrogen or halo; and "a", "b", "c", "d", "e", "f", and "h" are alternative sites for optional double bonds.

One class of preferred steroids has "b" as
 20 a double bond, particularly wherein "e" or "d" is also a double bond. Another preferred class has "a" and "c" as double bonds. Yet another preferred class contains P₃ as a methyl group, "h" as an optional double bond, and P₅ as methylene or one or two

Figs. 8C & D: Electrodermal activity was measured as described. Changes (measured in $x\Omega$) in response due to delivery of vomeropherins to the VNO of each subject are shown in male (C) and female (D) subjects.

Figs. 8E & F: Alpha-cortical activity was measured as described. Changes in response due to delivery of vomeropherins to the VNO of male (E) and female (F) subjects.

10 Figs. 8G & H: Skin temperature (ST) was measured as described. Changes in response due to delivery of vomeropherins to the VNO of each subject are shown in male (G) and female (H) subjects.

The Compounds in the graphs are:

- 15 A = 1, 3, 5(10),16-Estratetraen-3-yl acetate
B = Androsta-4,16-dien-3-one
C = 1,3,5(10),16-Estratetraen-3-ol
D = 3-Methoxy-Estra-1,3,5(10),16-tetraene
E = Androsta-4,16-dien-3 α -ol
20 F = Androsta- 4,16-dien-3 β -ol

Figure 9 depicts electro-olfactograms of male and female subjects induced by stimulation of the OE with olfactants and vomeropherins A: 400 fmoles of the olfactants l-carvone and cineole as well as 200 fmoles of the vomeropherins A, B, C, D and F; and the stereoisomer E were applied separately as one second pulses to the OE of 20 subjects (both male and female) and each EOG response was recorded as described. The olfactants as well as E and B produced significant ($p < 0.01$) local response. B: 400 fmoles of the olfactants l-carvone and cineole do not

resentfulness, bitterness, timidity, laziness and the like.

"Androstane steroids" are aliphatic polycyclic hydrocarbons characterized by a four-ring steroidal structure with a methylation at the 10- and 13-positions. An Androstene is a subset of Androstanes commonly understood to mean that the compound has at least one double bond. Commonly, unless a compound is described as a gonane, it is understood that the compound has a 19- carbon group. Furthermore, all derivatives which have the structural characteristics described above are also referred to generically as Androstane steroids.

A "chemoreceptor" is a receptor molecule displayed on the surface of a "chemosensory" neuroepithelial cell which binds in a stereospecific fashion to a particular ligand or ligands. This specific binding initiates a signal transduction which initiates an afferent nerve impulse. Chemoreceptors are found, inter alia, in taste buds, olfactory epithelium and vomeronasal tissue.

"Estrene steroids", as the term is used herein, are aliphatic polycyclic hydrocarbons with a four-ring steroidal structure, at least one double bond in the A-ring, no methylation at the 10-position and an oxo, hydroxyl or hydroxyl derivative such as an alkoxy, ester, benzoate, cypionate, sulfate or glucuronide, at the 3-position. Derivatives which contain these structural characteristics are also referred to generically as Estrene steroids.

The following structure shows the four-ring steroidal structure common to Androstane and Estrene steroids. In describing the location of groups and

alkoxymethyl, lower alkyl, hydroxyalkyl,
acyloxyalkyl, and alkoxylalkyl; P₄ is selected from
the group consisting of hydrogen, oxo, halo, hydroxy,
alkoxy, and acyloxy; P₅ represents one or 2
5 substituents, wherein P₅ comprises one or two
hydrogen atoms, methyl, methylene, or one or two halo
atoms; P₆ is hydrogen or halo; and "a", "b", "c",
"d", "e", "f", and "h" are alternative sites for
optional double bonds.

10 2. A composition according to Claim 1
wherein P₃ a methyl or other lower alkyl.

 3. A composition according to Claim 1
wherein "b" is a double bond.

 4. A composition according to Claim 3
15 wherein P₁ is oxo.

 5. A composition according to Claim 4
wherein P₄ is hydrogen.

 6. A composition according to Claim 5
wherein said steroid is selected from the group
20 consisting of androsta-4,16-dien-3-one, androst-4-en-
3-one, 17-methylene-androst-4-en-3-one, 18-nor-17-

methyl-androsta-4,17(13)-dien-3-one, 19-hydroxy-androsta-4,16-dien-3-one, 19-hydroxy-androst-4-en-3-one, 19-hydroxy-17-methylene-androst-4-en-3-one and 18-nor-17-methyl-19-hydroxy-androsta-4,17(13)-dien-3-one.

7. A composition according to Claim 4 wherein P_4 is oxo.

8. A composition according to Claim 7 selected from the group consisting of androsta-4,16-dien-3,6-dione, androst-4-en-3,6-dione, 17-methylene-androst-4-en-3,6-dione and 18-nor-17-methyl-androsta-4,17(13)-dien-3,6-dione.

9. A composition according to Claim 4 wherein P_4 is hydroxy.

10. A composition according to Claim 9 wherein said steroid is selected from the group consisting of 6α -hydroxy-androsta-4,16-dien-3-one, 6α -hydroxy-androst-4-en-3-one, 17-methylene- 6α -hydroxy-androst-4-en-3-one and 18-nor-17-methyl- 6α -hydroxy-androsta-4,17(13)-dien-3-one.

11. A composition according to Claim 3 wherein P_1 is hydroxy.

12. A composition according to Claim 11 wherein said steroid is selected from the group
5 consisting of androsta-4,16-dien-3 α -ol, androsta-4,16-dien-3 β -ol, androst-4-en-3 α -ol, androst-4-en-3 β -ol, 17-methylene-androst-4-en-3 α -ol, 17-methylene-androst-4-en-3 β -ol, 18-nor-17-methyl-androsta-4,17(13)-dien-3 α -ol, 18-nor-17-methyl-androsta-4,
10 17(13)-dien-3 β -ol.

13. A composition according to Claim 3 wherein "e" or "d" is a double bond.

14. A composition according to Claim 13 wherein said steroid is selected from the group
15 consisting of androsta-4, 6,16-trien-3-one, androsta-4,6-dien-3-one, 17-methylene-androsta-4,16-dien-3-one, 18-nor-17-methyl-androsta-4,6,17(13)-trien-3-one, androsta-1,4,16-trien-3-one, androsta-1,4-dien-3-one, 17-methylene-1,4-dien-3-one, 18-nor-17-methyl-
20 androsta-1,4,17(13)-trien-3-one, androsta-1,4,16-trien-3 α -ol, androsta-1,4-dien-3 α -ol, 17-methylene-androsta-1,4-dien-3 α -ol, 18-nor-17-methyl-androsta-1,4,17(13)-trien-3 α -ol.

15. A composition according to Claim 1 wherein "c" is a double bond.

16. A composition according to Claim 15 wherein said steroid is selected from the group
5 consisting of androsta-5(6),16-dien-3 α -ol, androst-5(6)-en-3 α -ol, 17-methylene-androst-5(6)-en-3 α -ol, 18-nor-17-methyl-androsta-5(6),17(13)-dien-3 α -ol.

17. A composition according to Claim 15 wherein "a" is a double bond.

10 18. A composition according to Claim 17 wherein said steroid is selected from the group consisting of 3-methoxy-androsta-3,5(6),16-triene, 3-methoxy-androsta-3,5(6)-diene, 17-methylene-androsta-3,5(6)-diene, 18-nor-17-methyl-3-methoxy-androsta-
15 3,5(6),17(13)-triene.

19. A composition according to Claim 1 wherein P₃ is methyl.

20. A composition according to Claim 19 wherein P₅ is methylene.

21. A composition according to Claim 20 wherein said steroid is selected from the group consisting of 17-methylene-androst-4-en-3-one, 17-methylene-androst-5(6)-en-3 α -ol, 17-methylene-
5 androst-4-en-3 α -ol, 17-methylene-androst-4-en-3 β -ol, 17-methylene-androsta-4,6-dien-3-one, 17-methylene-androst-4-en-3,6-dione, 17-methylene-19-hydroxy-androst-4-en-3-one, 17-methylene-3-methoxy-androsta-3,5(6)-diene, 17-methylene-androsta-1,4-dien-3-one,
10 17-methylene-androsta-1,4-dien-3 α -ol, 17-methylene-6 α -hydroxy-androst-4-en-3-one.

22. A composition according to Claim 19 wherein P₅ is two hydrogen atoms.

23. A composition according to Claim 22
15 wherein said steroid is selected from the group consisting of androst-4-en-3-one, androst-5(6)-en-3 α -ol, androst-4-en-3 α -ol, androst-4-en-3 β -ol, androsta-4,6-dien-3-one, androst-4-en-3,6-dione, 19-hydroxy-androst-4-en-3-one, 3-methoxy-androsta-3,5(6)-diene,
20 androsta-1,4-dien-3-one, androsta-1,4-dien-3 α -ol, androst-4-en-6 α -ol-3-one.

24. A composition according to Claim 1 wherein P₃ is methyl and "h" is a double bond.

25. A composition according to Claim 24 wherein said steroid is selected from the group consisting of androsta-4,16-dien-3-one, 3 α -hydroxy-androsta-5(6),16-diene, 3 α -hydroxy-androsta-4,16-
5 diene, 3 β -hydroxy-androsta-4,16-diene, androsta-4,6,16-trien-3-one, androsta-4,16-dien-3,6-dione, 19-hydroxy-androsta-4,16-dien-3-one, 3-methoxy-androsta-3,5(6),16-triene, androsta-1,4,16-trien-3-one, 3 α -hydroxy-androsta-1,4,16-triene, androsta-4,16-dien-
10 6 α -ol-3-one.

26. A composition according to Claim 1 wherein P₅ is methyl and "f" is a double bond.

27. A composition according to Claim 26 wherein said steroid is selected from the group
15 consisteing of 18-nor-17-methyl-androsta-4,17(13)-dien-3-one, 18-nor-17-methyl-androsta-5(6),17(13)-dien-3 α -ol, 18-nor-17-methyl-androsta-4,17(13)-dien-3 α -ol, 18-nor-17-methyl-androsta-4,17(13)-dien-3 β -ol, 18-nor-17-methyl-androsta-4,6,17(13)-trien-3-one, 18-
20 nor-17-methyl-androsta-4,17(13)-dien-3,6-dione, 18-nor-17-methyl-19-hydroxy-androsta-4,17(13)-dien-3-one, 18-nor-17-methyl-3-methoxy-androsta-3,5(6),17(13)-triene, 18-nor-17-methyl-androsta-1,4,17(13)-trien-3-one, 18-nor-17-methyl-androsta-

1,4-dien-3 α -ol and 18-nor-17-methyl-androsta-
4,17(13)-dien-6 α -ol-3-one.

28. The pharmaceutical composition of any
of Claims 1 through 27 wherein said steroid is
5 dissolved in said carrier.

29. The pharmaceutical composition of any
of Claims 1 through 27 wherein said composition is in
a liquid form.

30. The pharmaceutical composition of any
10 of Claims 1 through 27 wherein said composition
further contains a pharmaceutically acceptable
ointment base.

31. The pharmaceutical composition of any
of Claims 1 through 27 which contains no more than
15 one of said steroids.

32. The pharmaceutical composition of any
of Claims 1 through 27 which contains more than one
of said steroids.

33. The pharmaceutical composition of any
of Claims 1 through 27 additionally comprising an
Estrene steroid.

34. A method of altering a hypothalamic
5 function of an individual, said method comprising:

providing a ligand for a chemoreceptor
displayed on the surface of nasal
neuroepithelial cell of said individual
wherein said cell is a part of tissue other
10 than olfactory epithelia; and,

administering said ligand within a nasal
passage of said individual such that said
ligand binds specifically to said receptor
and results in an alteration of
15 hypothalamic function of said individual.

35. A method of altering an autonomic
function of an individual, said method comprising:


providing a ligand for a chemoreceptor of a
nasal neuroepithelial cell of-said
20 individual wherein said cell is a part of
tissue other than olfactory epithelia; and,

administering said ligand within a nasal
passage of said individual such that said
ligand binds specifically to said receptor
and results in an alteration of
5 hypothalamic function of said individual.

36. The method of Claim 35 wherein said
neuroepithelial cell is located within a vomeronasal
organ of said individual.

37. The method of claim 36 wherein said
10 ligand comprises an Androstane.

38. The method of claim 37 wherein said
Androstane has the formula:



wherein P_1 is selected from the group consisting of oxo, α -(β -) hydroxy, α -(β -) acetoxy, α -(β -) propionyxy, α -(β -) methoxy, α -(β -) lower acyloxy, α -(β -) lower alkyloxy, and α -(β -) benzoyloxy; P_2 is
5 selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl, alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P_3 is absent or is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl,
10 alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P_4 is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P_5 represents one or 2 substituents, wherein P_5 comprises one or two
15 hydrogen atoms, methyl, methylene, or one or two halo atoms; P_6 is hydrogen or halo; and "a", "b", "c", "d", "e", "f", and "h" are alternative sites for optional double bonds.

39. A method according to Claim 38 wherein
20 P_3 a methyl or other lower alkyl.

40. A method according to Claim 38 wherein
"b" is a double bond.

41. A method according to Claim 40 wherein P_1 is oxo.

42. A method according to Claim 41 wherein P_4 is hydrogen.

5 43. A method according to Claim 42 wherein said steroid is selected from the group consisting of androsta-4,16-dien-3-one, androst-4-en-3-one, 17-methylene-androst-4-en-3-one, 18-nor-17-methyl-androsta-4,17(13)-dien-3-one, 19-hydroxy-androsta-
10 4,16-dien-3-one, 19-hydroxy-androst-4-en-3-one, 19-hydroxy-17-methylene-androst-4-en-3-one and 18-nor-17-methyl-19-hydroxy-androsta-4,17(13)-dien-3-one.

44. A method according to Claim 41 wherein P_4 is oxo.

15 45. A method according to Claim 44 selected from the group consisting of androsta-4,16-dien-3,6-dione, androst-4-en-3,6-dione, 17-methylene-androst-4-en-3,6-dione and 18-nor-17-methyl-androsta-4,17(13)-dien-3,6-dione.

20 46. A method according to Claim 41 wherein P_4 is hydroxy.

47. A method according to Claim 46 wherein said steroid is selected from the group consisting of 6 α -hydroxy-androsta-4,16-dien-3-one, 6 α -hydroxy-androst-4-en-3-one, 17-methylene-6 α -hydroxy-androst-4-en-3-one and 18-nor-17-methyl-6 α -hydroxy-androsta-4,17(13)-dien-3-one.

48. A method according to Claim 40 wherein P₁ is hydroxy.

49. A method according to Claim 48 wherein said steroid is selected from the group consisting of androsta-4,16-dien-3 α -ol, androsta-4,16-dien-3 β -ol, androst-4-en-3 α -ol, androst-4-en-3 β -ol, 17-methylene-androst-4-en-3 α -ol, 17-methylene-androst-4-en-3 β -ol, 18-nor-17-methyl-androsta-4,17(13)-dien-3 α -ol, 18-nor-17-methyl-androsta-4,17(13)-dien-3 β -ol.

50. A method according to Claim 40 wherein "e" or "d" is a double bond.

51. A method according to Claim 50 wherein said steroid is selected from the group consisting of androsta-4,6,16-trien-3-one, androsta-4,6-dien-3-one, 17-methylene-androsta-4,16-dien-3-one, 18-nor-17-methyl-androsta-4,6,17(13)-trien-3-one, androsta-

1,4,16-trien-3-one, androsta-1,4-dien-3-one, 17-methylene-1,4-dien-3-one, 18-nor-17-methyl-androsta-1,4,17(13)-trien-3-one, androsta-1,4,16-trien-3 α -ol, androsta-1,4-dien-3 α -ol, 17-methylene-androsta-1,4-
5 dien-3 α -ol, 18-nor-17-methyl-androsta-1, 4,17(13)-trien-3 α -ol.

52. A method according to Claim 38 wherein "c" is a double bond.

53. A method according to Claim 52 wherein
10 said steroid is selected from the group consisting of androsta-5(6),16-dien-3 α -ol, androst-5(6)-en-3 α -ol, 17-methylene-androst-5(6)-en-3 α -ol, 18-nor-17-methyl-androsta-5(6),17(13)-dien-3 α -ol.

54. A method according to Claim 38 wherein
15 "a" is a double bond.

55. A method according to Claim 54 wherein said steroid is selected from the group consisting of 3-methoxy-androsta-3,5(6),16-triene, 3-methoxy-androsta-3,5(6)-diene, 17-methylene-androsta-3,5(6)-
20 diene, 18-nor-17-methyl-3-methoxy-androsta-3,5(6),17(13)-triene.

56. A method according to Claim 38 wherein P_3 is methyl.

57. A method according to Claim 56 wherein P_5 is methylene.

5 58. A method according to Claim 57 wherein
said steroid is selected from the group consisting of
17-methylene-androst-4-en-3-one, 17-methylene-
androst-5(6)-en-3 α -ol, 17-methylene-androst-4-en-3 α -
ol, 17-methylene-androst-4-en-3 β -ol, 17-methylene-
10 androsta-4,6-dien-3-one, 17-methylene-androst-4-en-
3,6-dione, 17-methylene-19-hydroxy-androst-4-en-3-
one, 17-methylene-3-methoxy-androsta-3,5(6)-diene,
17-methylene-androsta-1,4-dien-3-one, 17-methylene-
androsta-1,4-dien-3 α -ol, 17-methylene-6 α -hydroxy-
15 androst-4-en-3-one.

59. A composition according to Claim 56
wherein P_5 is two hydrogen atoms.

60. A composition according to Claim 59
wherein said steroid is selected from the group
20 consisting of androst-4-en-3-one, androst-5(6)-en-3 α -
ol, androst-4-en-3 α -ol, androst-4-en-3 β -ol, androsta-
4,6-dien-3-one, androst-4-en-3,6-dione, 19-hydroxy-

androst-4-en-3-one, 3-methoxy-androsta-3,5(6)-diene,
androsta-1,4-dien-3-one, androsta-1,4-dien-3 α -ol,
androst-4-en-6 α -ol-3-one.

61. A method according to Claim 38 wherein
5 P₃ is methyl and "h" is a double bond.

62. A method according to Claim 61 wherein
said steroid is selected from the group consisting of
androsta-4,16-dien-3-one, 3 α -hydroxy-androsta-
5(6),16-diene, 3 α -hydroxy-androsta-4,16-diene, 3 β -
10 hydroxy-androsta-4,16-diene, androsta-4,6,16-trien-3-
one, androsta-4,16-dien-3,6-dione, 19-hydroxy-
androsta-4,16-dien-3-one, 3-methoxy-androsta-
3,5(6),16-triene, androsta-1,4,16-trien-3-one, 3 α -
hydroxy-androsta-1,4,16-triene, androsta-4,16-dien-
15 6 α -ol-3-one.

63. A method according to Claim 38 wherein
P₅ is methyl and "f" is a double bond.

64. A composition according to Claim 63
wherein said steroid is selected from the group
20 consisteing of 18-nor-17-methyl-androsta-4,17(13)-
dien-3-one, 18-nor-17-methyl-androsta-5(6),17(13)-
dien-3 α -ol, 18-nor-17-methyl-androsta-4,17(13)-dien-

3 α -ol, 18-nor-17-methyl-androsta-4,17(13)-dien-3 β -ol,
18-nor-17-methyl-androsta-4,6,17(13)-trien-3-one, 18-
nor-17-methyl-androsta-4,17(13)-dien-3,6-dione, 18-
nor-17-methyl-19-hydroxy-androsta-4,17(13)-dien-3-
5 one, 18-nor-17-methyl-3-methoxy-androsta-
3,5(6),17(13)-triene, 18-nor-17-methyl-androsta-
1,4,17(13)-trien-3-one, 18-nor-17-methyl-androsta-
1,4-dien-3 α -ol and 18-nor-17-methyl-androsta-
4,17(13)-dien-6 α -ol-3-one.

10 65. The method of any of Claims 34 through
64 wherein the amount of said ligand that is
administered is at least about 100 picograms, but no
more than about 100 micrograms.

15 66. The method of Claim 65 wherein the
amount of said ligand that is administered is at
least about 1 nanograms, but no more than about 10
micrograms.

20 67. The method of Claim 66 wherein the
amount of said ligand that is administered is at
least about 10 nanograms, but no more than about 1
microgram.

68. The method of any of Claims 34 through 64 further comprising preparing a pharmaceutical composition of said ligand dissolved in a pharmaceutically acceptable carrier.

5 69. The method of Claim 68 wherein said pharmaceutical composition is an ointment.

70. The method of Claim 68 wherein said pharmaceutical composition is liquid.

71. The method of Claim 68 wherein the
10 administration is by aerosol.

72. The method of any of Claims 34 through 64 wherein more than one Androstane steroid is administered.

73. The method of any of Claims 34 through
15 64 further comprising nasally co-administering to said individual an Estrene steroid.

74. The method of any of Claims 34 through 64 wherein said function is the diminution of negative affect.

75. The method of any of Claims 34 through 64 wherein said individual is a woman.

76. The method of Claim 75 wherein said alteration of hypothalamic function results in a
5 reduction of premenstrual stress in women.

77. The method of any of Claims 34 through 64 wherein said function is the alleviation of the symptoms of psychoses, depression or anxiety.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/09349

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : A61K 31/56

US CL : 514/170,177,178,182

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/170,177,178,182

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS and CAS ONLINE: androstane, psych?, depress?, anxiet?, hypothal?, autonom?, chemorecept?, steroid

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,E	US, A, 5,272,134 (Berliner) 21 December 1993, see entire document.	1-54
Y	US, A, 3,908,007 (Itil et al.) 23 September 1975, see column 1, line 4-column 2, line 2.	22-54
Y	US, A, 4,330,538 (Itil et al.) 18 May 1982, see abstract and column 1-column 2, line 14.	22-54



Further documents are listed in the continuation of Box C.



See patent family annex.

Special categories of cited documents:		T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A	document defining the general state of the art which is not considered to be part of particular relevance		
E	earlier document published on or after the international filing date	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O	document referring to an oral disclosure, use, exhibition or other means		
P	document published prior to the international filing date but later than the priority date claimed	Z	document member of the same patent family

Date of the actual completion of the international search

04 January 1993


Date of mailing of the international search report

27 JAN 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

KIMBERLY JORDAN



Telephone No. (703) 205-1235




FIG. 1.




FIG. 2





FIG. 3.






FIG 4B.




FIG 4C

4/10




FIG. 5A.

5/10




FIG. 5B.

6/10




FIG. 6A.




FIG. 6B.




FIG. 6C.




FIG. 6D.




FIG. 6E.




FIG. 6F.

7/10




FIG. 7.




FIG. 9A.




FIG. 9B.

8/10




FIG. 8A.




FIG. 8B.




FIG. 8C.




FIG. 8D.



FIG. 8E.




FIG. 8F.




FIG. 8G.




FIG. 8H.

9/10

FIG. 10A




FIG. 10B





FIG. 10C



10/10

FIG. IIA.




FIG. IIB.




FIG. IIC.

